

Approved

To learn more about how developmental improvements have been measured in babies fed **Enfamil[®] LIPIL[®]**, visit Enfamil.com/Infantment. If you choose to use infant formula, ask your baby's doctor about the many benefits of **Enfamil LIPIL**, our closest formula to breast milk.

Only Enfamil[®]. Because it's your baby.[™]

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula.
2. Pour desired amount of water into the bottle. Add powder.
3. Cap bottle and SHAKE WELL.

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 unpacked level scoop (8.5g)
4 fl oz bottle	4 fl oz	2 unpacked level scoops (17g)
8 fl oz bottle	8 fl oz	4 unpacked level scoops (34g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **USE BY DATE ON BOTTOM OF CAN**

We are here for you. Call us toll free at 1-800-BABY123, 8 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

Only Brand CLINICALLY PROVEN To Improve VISUAL & MENTAL DEVELOPMENT Improvements Shown Beyond Baby's First Year Up to 18 Months!

Enfamil

LIPIL[®]

A blend of DHA & ARA

Trusted by Moms & Pediatricians

Infant Formula
Milk-Based Powder
with Iron

12.9 OZ (366 g)

Babies 0-12 Months



NUTRIENTS (Normal Dilution)	Per 100 Calories (5 fl oz)	INGREDIENTS: NON-FAT MILK, LACTOSE, VEGETABLE OIL (PALM OLEIN, SOY COCOBUT AND HIGH OLEIC SUNFLOWER OILS), WHEY PROTEIN CONCENTRATE, AND LESS THAN 1%: MORTIERELLA ALPINA OIL*, CRYPTOCODINIUM CORNI OIL** VITAMIN K ₁ PALMITATE, VITAMIN D ₃ , VITAMINE ACETATE, VITAMIN D ₆ , VITAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B ₆ , HYDROXYLACETATE, VITAMIN K ₁ , THIAMIN HYDROCHLORIDE, VITAMIN B ₁₂ , NICOTINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLENE CHLORIDE, INOSITOL, CALCIUM CARBOXYMAGNESIUM OXIDE, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, COPPER SULFATE, POTASSIUM CHLORIDE, POTASSIUM CITRATE, POTASSIUM HYDROXIDE, SODIUM SELENITE, TURINIC, CARANTINE, NUCLEOTIDES (ADENSINE 5'-MONOPHOSPHATE, CYTIDINE 5'-MONOPHOSPHATE, DISODIUM GUANOSINE 5'-MONOPHOSPHATE, DISODIUM URIDINE 5'-MONOPHOSPHATE).
PROTEIN	g ... 2.1	
FAT	g ... 5.3	
CARBOHYDRATE	g ... 10.9	
WATER	g ... 134	
LINOLEIC ACID	mg ... 260	
VITAMINS		
A	IU ... 300	
D	IU ... 60	
E	IU ... 2	
K	mcg ... 8	
THIAMIN (B ₁)	mcg ... 80	
RIBOFLAVIN (B ₂)	mcg ... 140	
B ₆	mcg ... 60	
B ₁₂	mcg ... 0.3	
NIACIN	mcg ... 1000	
FOLIC ACID (FOLACIN)	mcg ... 16	
PANTOTHENIC ACID	mcg ... 500	
BIOTIN	mcg ... 3	
C (ASCORBIC ACID)	mg ... 12	
CHOLINE	mg ... 24	
INOSITOL	mg ... 6	
MINERALS		
CALCIUM	mg ... 78	
PHOSPHORUS	mg ... 43	
MAGNESIUM	mg ... 8	
IRON	mg ... 1.8	
ZINC	mg ... 1	
MANGANESE	mcg ... 15	
COPPER	mcg ... 75	
IODINE	mcg ... 10	
SELENIUM	mcg ... 2.8	
SODIUM	mg ... 27	
POTASSIUM CHLORIDE	mg ... 63	

WAL-MART
If this formula is available only from the Enfamil[®] brand, please call 1-800-WAL-MART



Not Approved

Growth

Proven healthy growth.
• Complete and balanced nutrition supported by over 100 years of experience

Brain & Eye

KQ scores & vision proven stronger in the breastfed by
• 17mg DHA and 34 mg ARA, per 100 Calories expert recommended levels

Immune System

Proven to help build a strong immune system
• Now with prebiotics similar to those found in breast milk

If you choose to use infant formula, ask your baby's doctor about new **Enfamil PREMIUM**.

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby. Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula.
2. Pour desired amount of water into the bottle. Add powder.
3. Cap bottle and SHAKE WELL.

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 unpacked level scoop (8.7g)
4 fl oz bottle	4 fl oz	2 unpacked level scoops (17.4g)
8 fl oz bottle	8 fl oz	4 unpacked level scoops (34.8g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **USE BY DATE ON BOTTOM OF CAN**

We are here for you. Call us toll free at 1-800-BABY123, 8 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

CLINICALLY PROVEN FOR HEALTHY DEVELOPMENT

Enfamil

PREMIUM

LIPIL[®]

NEW Clinically Proven Triple Health Guard Growth Brain & Eye Immune System

Recommended by Pediatricians & Trusted by Moms

Infant Formula
Milk-Based Powder with Iron

12.5 OZ (354g)



NUTRIENTS (Normal Dilution) per 100 Calories (5 fl oz)	INGREDIENTS: NON-FAT MILK, LACTOSE, VEGETABLE OIL (PALM OLEIN, SOY COCOBUT AND HIGH OLEIC SUNFLOWER OILS), WHEY PROTEIN CONCENTRATE, GALACTOOLIGOSACCHARIDES (GOS) [†] AND LESS THAN 1%: MORTIERELLA ALPINA OIL*, CRYPTOCODINIUM CORNI OIL** SOY LECITHIN, VITAMIN A PALMITATE, VITAMIN D ₃ , VITAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B ₆ , HYDROXYLACETATE, VITAMIN K ₁ , THIAMIN HYDROCHLORIDE, VITAMIN B ₁₂ , NICOTINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLENE CHLORIDE, INOSITOL, CALCIUM CARBOXYMAGNESIUM OXIDE, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, COPPER SULFATE, POTASSIUM CHLORIDE, POTASSIUM CITRATE, POTASSIUM HYDROXIDE, SODIUM SELENITE, TURINIC, CARANTINE, NUCLEOTIDES (ADENSINE 5'-MONOPHOSPHATE, CYTIDINE 5'-MONOPHOSPHATE, DISODIUM GUANOSINE 5'-MONOPHOSPHATE, DISODIUM URIDINE 5'-MONOPHOSPHATE).
PROTEIN	g ... 2.1
FAT	g ... 5.3
CARBOHYDRATE	g ... 11
WATER	g ... 133
LINOLEIC ACID	mg ... 260
VITAMINS	
A	IU ... 300
D	IU ... 60
E	IU ... 2
K	mcg ... 8
THIAMIN (B ₁)	mcg ... 80
RIBOFLAVIN (B ₂)	mcg ... 140
B ₆	mcg ... 60
B ₁₂	mcg ... 0.3
NIACIN	mcg ... 1000
FOLIC ACID (FOLACIN)	mcg ... 16
PANTOTHENIC ACID	mcg ... 500
BIOTIN	mcg ... 3
C (ASCORBIC ACID)	mg ... 12
CHOLINE	mg ... 24
INOSITOL	mg ... 6
MINERALS	
CALCIUM	mg ... 78
PHOSPHORUS	mg ... 43
MAGNESIUM	mg ... 8
IRON	mg ... 1.8
ZINC	mg ... 1
MANGANESE	mcg ... 15
COPPER	mcg ... 75
IODINE	mcg ... 10
SELENIUM	mcg ... 2.8
SODIUM	mg ... 27
POTASSIUM CHLORIDE	mg ... 63

This exclusive formula is available only from the Enfamil[®] brand



Nutramigen® LIPIL® with Enfloira™ LGG is a hypoallergenic, lactose-free formula for babies who develop cow's milk protein allergy. Additionally, Nutramigen LIPIL with Enfloira LGG includes Lactobacillus GG which has been shown to promote immune system balance. Experts agree on the many benefits of breast milk. If you choose to use infant formula, ask your baby's doctor about the many benefits of Nutramigen LIPIL with Enfloira LGG.

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who are immunocompromised unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula.
2. Pour desired amount of water into the bottle. Add powder.
3. Cap bottle and SHAKE WELL.

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 PACKED level scoop (9 g)
4 fl oz bottle	4 fl oz	2 PACKED level scoops (18 g)
8 fl oz bottle	8 fl oz	4 PACKED level scoops (36 g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not warm. Warming may limit benefits of LGG. Microwaving formula may cause serious burns to baby. Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly.

Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

The product does not look or taste like milk or soy formulas. It may separate in the refrigerator. Shake well before feeding. USE BY DATE ON BOTTOM OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 9 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

AS AN ADDED SAFETY PRECAUTION, PLEASE CHECK THAT "NUTRI GO" IS EMBOSSED ON THE BOTTOM OF THIS CAN

NEW FORMULATION!

Nutramigen™

LIPIL®

DHA (Omega-3) & ARA



For Babies with COLIC due to Cow's Milk Protein Allergy

Infant Formula
Hypoallergenic Powder with Iron



12.6 OZ (357 g)



NUTRIENTS (Normal Dilution) per 100 Calories (5 fl oz)	
Protein	g 2.8
Fat	g 5.3
Carbohydrate	g 10.3
Water	g 133
Linoleic acid	mg 860
VITAMINS	
A	IU 300
D	IU 50
E	IU 2
K	mcg 9
Thiamin (B1)	mcg 80
Riboflavin (B2)	mcg 90
B6	mcg 60
B12	mcg 0.3
Niacin	mcg 1000
Folic acid (Folacin)	mcg 16
Pantothenic acid	mcg 500
Biotin	mcg 3
C (Ascorbic acid)	mg 12
Choline	mg 24
Inositol	mg 17
MINERALS	
Calcium	mg 94
Phosphorus	mg 52
Magnesium	mg 8
Iron	mg 1.8
Zinc	mg 1
Manganese	mcg 25
Copper	mcg 75
Iodine	mcg 15
Selenium	mcg 2.8
Sodium	mg 47
Potassium	mg 110
Chloride	mg 86

INGREDIENTS: CORN SYRUP SOLIDS (46%), VEGETABLE OIL (PALM OIL, SOY, COCOBUT, AND HIGH OLEIC SUNFLOWER OILS) (25%), CASEIN HYDROLYSATE* (FROM MILK) (17%), MODIFIED CORN STARCH (7%) AND LESS THAN 2% MOIFRELLA ALPINA OIL** CRYPTOCODONUM CORNUI OIL† ACETYLATED MONOGLYCERIDES, VITAMIN A PALMITATE, VITAMIN D3, VITAMIN E ACETATE, VITAMIN K1, THIAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B6 HYDROCHLORIDE, VITAMIN B12, NIACINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLINE CHLORIDE, INOSITOL, CALCIUM CITRATE, CALCIUM HYDROXIDE, CALCIUM PHOSPHATE, MAGNESIUM OXIDE, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, COPPER SULFATE, SODIUM IODIDE, SODIUM SELENITE, SODIUM CITRATE, POTASSIUM CITRATE, POTASSIUM CHLORIDE, L-CYSTEINE, L-TYROSINE, L-TRYPTOPHAN, TAURINE, L-CARNITINE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47721 U.S.A.
©2009 Mead Johnson & Company

*MODIFIED TO BE BETTER TOLERATED IN MILK-ALLERGIC BABIES
**A SOURCE OF ARACHIDONIC ACID (ARA)
†A SOURCE OF DOCOSAHEXAENOIC ACID (DHA)
LGG is a registered trademark of Valio Ltd.
Filled by weight, not by volume; some settling may occur.
Makes 67 fl oz

This exclusive formula is available only from the Enfamil® brand.



Nutramigen® LIPIL® is a hypoallergenic, lactose-free formula for babies who develop cow's milk protein allergy symptoms such as excessive crying or rash. LIPIL is a blend of DHA and ARA, important nutrients also found in breast milk that promote brain and eye development. Experts agree on the many benefits of breast milk. If you choose to use infant formula, ask your baby's doctor about the many benefits of Nutramigen LIPIL.

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula.
2. Pour desired amount of water into the bottle. Add powder.
3. Cap bottle and SHAKE WELL.

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 PACKED level scoop (9 g)
4 fl oz bottle	4 fl oz	2 PACKED level scoops (18 g)
8 fl oz bottle	8 fl oz	4 PACKED level scoops (36 g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Nutramigen LIPIL does not look or taste like milk or soy formulas. It may separate in the refrigerator. Shake well before feeding. USE BY DATE ON BOTTOM OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 9 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

Nutramigen™

LIPIL®

A blend of DHA & ARA—
Nutrients Found in Breast Milk

For Babies with COLIC due to Cow's Milk Protein Allergy

Hypoallergenic
Infant Formula Powder
Iron Fortified



1 LB (454 g) 16 OZ Babies 0-12 Months



NUTRIENTS (Normal Dilution) Per 100 Calories (5 fl oz)	
Protein	g 2.8
Fat	g 5.3
Carbohydrate	g 10.3
Water	g 133
Linoleic acid	mg 860
Vitamins:	
A	IU 300
D	IU 50
E	IU 2
K	mcg 9
Thiamin (Vitamin B1)	mcg 80
Riboflavin (Vitamin B2)	mcg 90
B6	mcg 60
B12	mcg 0.3
Niacin	mcg 1000
Folic acid (Folacin)	mcg 16
Pantothenic acid	mcg 500
Biotin	mcg 3
C (Ascorbic acid)	mg 12
Choline	mg 24
Inositol	mg 17
Minerals:	
Calcium	mg 94
Phosphorus	mg 52
Magnesium	mg 11
Iron	mg 1.8
Zinc	mg 1
Manganese	mcg 25
Copper	mcg 75
Iodine	mcg 15
Selenium	mcg 2.8
Sodium	mg 47
Potassium	mg 110
Chloride	mg 86

INGREDIENTS: CORN SYRUP SOLIDS (46%), VEGETABLE OIL (PALM OIL, SOY, COCOBUT, AND HIGH OLEIC SUNFLOWER OILS) (25%), CASEIN HYDROLYSATE* (FROM MILK) (17%), MODIFIED CORN STARCH (7%) AND LESS THAN 2% MOIFRELLA ALPINA OIL** CRYPTOCODONUM CORNUI OIL† ACETYLATED MONOGLYCERIDES, VITAMIN A PALMITATE, VITAMIN D3, VITAMIN E ACETATE, VITAMIN K1, THIAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B6 HYDROCHLORIDE, VITAMIN B12, NIACINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLINE CHLORIDE, INOSITOL, CALCIUM CITRATE, CALCIUM HYDROXIDE, CALCIUM PHOSPHATE, MAGNESIUM OXIDE, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, COPPER SULFATE, SODIUM IODIDE, SODIUM SELENITE, SODIUM CITRATE, POTASSIUM CITRATE, POTASSIUM CHLORIDE, L-CYSTEINE, L-TYROSINE, L-TRYPTOPHAN, TAURINE, L-CARNITINE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47721 U.S.A.
©2009 Mead Johnson & Company

*MODIFIED TO BE BETTER TOLERATED IN MILK-ALLERGIC BABIES
**A SOURCE OF ARACHIDONIC ACID (ARA)
†A SOURCE OF DOCOSAHEXAENOIC ACID (DHA)
Filled by weight, not by volume; some settling may occur.
Makes 131 fl oz

This scientifically developed formula is not available as a store brand.



Enfamil® Gentlease® LIPIL® has a unique design for babies with fussiness or gas. It is an easy-to-digest formula that contains partially broken down whey and casein milk proteins (proteins found in breast milk) and reduced lactose (about 1/3 the lactose of a full-lactose, routine, milk-based formula). LIPIL is a blend of DHA and ARA, important nutrients also found in breast milk that promote brain and eye development. Experts agree on the many benefits of breast milk. If you choose to use infant formula, ask your baby's doctor about the many benefits of Enfamil Gentlease LIPIL.

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula. 
2. Pour desired amount of water into the bottle. Add powder. 
3. Cap bottle and SHAKE WELL. 

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 unpacked level scoop (8.7 g)
4 fl oz bottle	4 fl oz	2 unpacked level scoops (17.4 g)
8 fl oz bottle	8 fl oz	4 unpacked level scoops (34.8 g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

USE BY DATE ON BOTTOM OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 9 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

NEW LOOK!

Enfamil™

Gentlease®

LIPIL®

DHA (Omega-3) & ARA

OUR #1 Choice for Fussiness and Gas

For Babies with Fussiness & Gas

Infant Formula
Milk-Based Powder with Iron - Partially Broken Down Proteins

12 OZ (340 g)



Babies 0-12 Months

NUTRIENTS (Normal Dilution)	Per 100 Calories (5 fl oz)
PROTEIN	g 2.3
FAT	g 5.3
CARBOHYDRATE	g 10.8
WATER	g 134
LINOLEIC ACID	mg 860

VITAMINS

A	IU 300
D	IU 60
E	IU 2
K	mcg 9
THIAMIN (B1)	mcg 80
RIBOFLAVIN (B2)	mcg 140
B6	mcg 60
B12	mcg 0.3
NIACIN	mcg 1000
FOLIC ACID (FOLACIN)	mcg 16
PANTOTHENIC ACID	mcg 500
BIOTIN	mcg 3
C (ASCORBIC ACID)	mg 12
CHOLINE	mg 24
INOSITOL	mg 6

MINERALS

CALCIUM	mg 82
PHOSPHORUS	mg 46
MAGNESIUM	mg 8
IRON	mg 1.8
ZINC	mg 1
MANGANESE	mcg 15
COPPER	mcg 75
IODINE	mcg 15
SELENIUM	mcg 2.8
SODIUM	mg 36
POTASSIUM CHLORIDE	mg 108

INGREDIENTS: CORN SYRUP SOLIDS, PARTIALLY HYDROLYZED NONFAT MILK AND WHEY PROTEIN CONCENTRATE SOLIDS (SOY), VEGETABLE OIL (PALM OLEIN, SOY, COCONUT) AND HIGH OLEIC SUNFLOWER OILS, AND LESS THAN 2% MORTIERELLA ALPINA OIL, CRYPTOCODINIUM COHNI OIL**, VITAMIN A, PALMITATE, VITAMIN D2, VITAMIN E ACETATE, VITAMIN K1, THIAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B6, HYDROCHLORIDE, VITAMIN B12, NIACINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLINE CHLORIDE, INOSITOL, CALCIUM CARBONATE, CALCIUM PHOSPHATE, MAGNESIUM PHOSPHATE, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, CUPRIC SULFATE, SODIUM SELENITE, SODIUM CITRATE, SODIUM HYDROXIDE, POTASSIUM HYDROXIDE, POTASSIUM CHLORIDE, TAURINE, AND L-CARNITINE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47721 U.S.A.
©2009 Mead Johnson & Company
*A SOURCE OF ARACHIDONIC ACID (ARA)
**A SOURCE OF DOCOSAHEXAENOIC ACID (DHA)

This exclusive formula is available only from the Enfamil® brand.



3 0087 8693 42 8

Enfamil A.R.® LIPIL® was specially designed for babies who spit up frequently. It contains an easy-to-digest rice starch designed to become thicker in the stomach and to help babies keep formula down gently. LIPIL is a blend of DHA and ARA, important nutrients also found in breast milk that promote brain and eye development. Experts agree on the many benefits of breast milk. If you choose to use infant formula, ask your baby's doctor about the many benefits of Enfamil A.R. LIPIL.

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula. 
2. Pour desired amount of water into the bottle. Add powder. 
3. Cap bottle and SHAKE WELL. Let bottle sit 5 minutes. 

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 unpacked level scoop (8.7 g)
4 fl oz bottle	4 fl oz	2 unpacked level scoops (17.4 g)
8 fl oz bottle	8 fl oz	4 unpacked level scoops (34.8 g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

USE BY DATE ON BOTTOM OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 9 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

NEW LOOK!

Enfamil™

A.R.®

LIPIL®

DHA (Omega-3) & ARA

Clinically Proven

For Babies Who Spit Up Frequently

Infant Formula
Milk-Based Powder with Iron - Thickened with Added Rice Starch

12.9 OZ (366 g)



Babies 0-12 Months

NUTRIENTS (Normal Dilution); per 100 Calories (5 fl oz)	
PROTEIN	g 2.5
FAT	g 5.1
CARBOHYDRATE	g 11
WATER	g 134
LINOLEIC ACID	mg 860

VITAMINS

A	IU 300
D	IU 60
E	IU 2
K	mcg 8
THIAMIN (B1)	mcg 80
RIBOFLAVIN (B2)	mcg 140
B6	mcg 60
B12	mcg 0.3
NIACIN	mcg 1000
FOLIC ACID (FOLACIN)	mcg 16
PANTOTHENIC ACID	mcg 500
BIOTIN	mcg 3
C (ASCORBIC ACID)	mg 12
CHOLINE	mg 24
INOSITOL	mg 6

MINERALS

CALCIUM	mg 78
PHOSPHORUS	mg 53
MAGNESIUM	mg 8
IRON	mg 1.8
ZINC	mg 1
MANGANESE	mcg 15
COPPER	mcg 75
IODINE	mcg 10
SELENIUM	mcg 2.8
SODIUM	mg 40
POTASSIUM CHLORIDE	mg 108

INGREDIENTS: NONFAT MILK, VEGETABLE OIL (PALM OLEIN, SOY, COCONUT) AND HIGH OLEIC SUNFLOWER OILS, RICE STARCH, LACTOSE, MALTODEXTRIN, AND LESS THAN 2% MORTIERELLA ALPINA OIL**, CRYPTOCODINIUM COHNI OIL**, VITAMIN A, PALMITATE, VITAMIN D2, VITAMIN E ACETATE, VITAMIN K1, THIAMIN HYDROCHLORIDE, RIBOFLAVIN, VITAMIN B6, HYDROCHLORIDE, VITAMIN B12, NIACINAMIDE, FOLIC ACID, CALCIUM CARBONATE, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLINE CHLORIDE, INOSITOL, FERROUS SULFATE, ZINC SULFATE, MANGANESE SULFATE, CUPRIC SULFATE, SODIUM SELENITE, SODIUM CITRATE, POTASSIUM HYDROXIDE, TAURINE, L-CARNITINE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47721 U.S.A.
©2009 Mead Johnson & Company
*A SOURCE OF ARACHIDONIC ACID (ARA)
**A SOURCE OF DOCOSAHEXAENOIC ACID (DHA)

This exclusive formula is available only from the Enfamil® brand.



3 0087 0201 42 3

Enfamil® EnfaCare® LIPIL®, when used in a program with other Enfamil LIPIL formulas, has been clinically proven to result in catch-up growth to the full-term, breast-fed baby. If you choose to use infant formula, ask your baby's doctor about the many benefits of EnfaCare LIPIL.

INSTRUCTIONS FOR PREPARATION & USE

WARNING: Your baby's health depends on carefully following the instructions below. Use only as directed by a medical professional. Improper hygiene, preparation, dilution, use or storage may result in severe harm. Although this powder is formulated for premature infants, powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula. 
2. Pour desired amount of water into the bottle. Add powder. 
3. Cap bottle and SHAKE WELL. 

Use the chart below for correct amounts of water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

To Make*	Water	Powder
2 fl oz bottle	2 fl oz	1 unpacked level scoop (9.8 g)
4 fl oz bottle	4 fl oz	2 unpacked level scoops (19.6 g)
8 fl oz bottle	8 fl oz	4 unpacked level scoops (39.2 g)

*Each scoop adds about 0.2 fl oz to the amount of prepared formula.

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula.

After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

USE BY DATE ON BOTTOM OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 8 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

NEW LOOK!

Enfamil

EnfaCare
LIPIL

DHA (Omega-3) & ARA

Clinically Proven Benefits

For Babies Who Were Born Prematurely

Infant Formula

Milk-Based Powder • 22 Cal/fl oz • with Iron

12.8 OZ (363 g)



NUTRIENTS
(Normal Dilution) per 100 Calories (4.5 fl oz)

PROTEIN	g	2.8
FAT	g	5.3
CARBOHYDRATE	g	10.4
WATER	g	117
LINOLEIC ACID	mg	960

VITAMINS

A	IU	450
D	IU	70
E	IU	4
K	mcg	8
THIAMIN (B1)	mcg	200
RIBOFLAVIN (B2)	mcg	200
B6	mcg	60
B12	mcg	0.3
NIACIN	mcg	1000
FOLIC ACID (FOLACIN)	mcg	26
PANTOTHENIC ACID	mcg	860
BIOTIN	mcg	5
C (ASCORBIC ACID)	mg	16
CHOLINE	mg	24
INOSITOL	mg	30

INGREDIENTS: NONFAT MILK, WHEY PROTEIN CONCENTRATE, COGNAC SYRUP SOLIDS, LACTOSE, HIGH OLEIC VEGETABLE OIL (SUNFLOWER AND/OR SAFFLOWER OIL), SOY OIL, MEDIUM CHAIN TRIGLYCERIDES (MCT OIL), COCONUT OIL, AND LESS THAN 2%: MORTIERELLA ALPINA OIL*, CRYPTOCODINIUM COINUI OIL**, VITAMIN A PALMITATE, VITAMIN D3, VITAMIN E ACETATE, VITAMIN K1, THIAMIN HYDROCHLORIDE, RIBOFLAVIN, NIACIN, HYDROCHLORIDE, VITAMIN B6, NIACINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLENE CHLORIDE, INOSITOL, CALCIUM CARBOXYLATE, CALCIUM PHOSPHATE, MANGANESE SULFATE, COPPER SULFATE, POTASSIUM IODIDE, SODIUM SELENITE, SODIUM CHLORIDE, POTASSIUM CITRATE, POTASSIUM HYDROXIDE, TAURINE, L-CARNITINE, NUCLEOTIDES: URIDINE, 5'-MONOPHOSPHATE, CYTIDINE, 5'-MONOPHOSPHATE, DISODIUM GUANOSINE, 5'-MONOPHOSPHATE, DISODIUM URIDINE, 5'-MONOPHOSPHATE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47711 U.S.A.
©2009 Mead Johnson & Company

*A SOURCE OF ARACHIDONIC ACID (ARA)
**A SOURCE OF DOCOSAHENAIC ACID (DHA)

Filled by weight, not by volume; some settling may occur. Makes 62 fl oz.



Enfamil® EnfaCare® LIPIL®, when used in a program with other Enfamil LIPIL formulas, has been clinically proven to result in catch-up growth to the full-term, breast-fed baby. If you choose to use infant formula, ask your baby's doctor about the many benefits of EnfaCare LIPIL.

INSTRUCTIONS FOR USE
Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Ask your baby's doctor about the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing feeding bottle. 
2. Clean can lid, SHAKE CAN WELL, and open. 
3. Pour into bottle(s). 

Warning: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened cans and prepared bottles can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 48 hours. Do not use opened can and/or prepared bottle if they are unrefrigerated for more than a total of 2 hours. Do not freeze prepared bottle.

After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

Storage: Store unopened cans at room temperature. Avoid excessive heat. Do not freeze. Use only as recommended by your baby's doctor for conditions such as prematurity.

USE BY DATE ON TOP OF CAN

We are here for you. Call us toll free at 1-800-BABY123, 8 AM to 4:30 PM Monday-Saturday, Central Time. Or visit our web site at www.enfamil.com

NEW LOOK!

Enfamil

EnfaCare
LIPIL

DHA (Omega-3) & ARA

Clinically Proven Benefits

For Babies Who Were Born Prematurely

Infant Formula

Milk-Based • 22 Cal/fl oz • with Iron

Ready to Use
Do Not Add Water

1 QT (946 mL)



NUTRIENTS per 100 Calories (4.5 fl oz)

PROTEIN	g	2.8
FAT	g	5.3
CARBOHYDRATE	g	10.4
WATER	g	117
LINOLEIC ACID	mg	960

VITAMINS

A	IU	450
D	IU	70
E	IU	4
K	mcg	8
THIAMIN (B1)	mcg	200
RIBOFLAVIN (B2)	mcg	200
B6	mcg	60
B12	mcg	0.3
NIACIN	mcg	1000
FOLIC ACID (FOLACIN)	mcg	26
PANTOTHENIC ACID	mcg	860
BIOTIN	mcg	5
C (ASCORBIC ACID)	mg	16
CHOLINE	mg	24
INOSITOL	mg	30

MINERALS

CALCIUM	mg	120
PHOSPHORUS	mg	66
MAGNESIUM	mg	8
IRON	mg	1.8
ZINC	mg	1.5
MANGANESE	mg	15
COPPER	mcg	629
IODINE	mcg	21
SELENIUM	mcg	2.8
SODIUM	mg	37
POTASSIUM	mg	105
CHLORIDE	mg	78

INGREDIENTS: WATER, NONFAT MILK, ANHYDROUS LACTOSE, WHEY PROTEIN CONCENTRATE, HIGH OLEIC VEGETABLE OIL (SUNFLOWER AND/OR SAFFLOWER OIL), SOY OIL, MEDIUM CHAIN TRIGLYCERIDES (MCT OIL), COCONUT OIL, AND LESS THAN 2%: MORTIERELLA ALPINA OIL*, CRYPTOCODINIUM COINUI OIL**, NIACIN, NIACINAMIDE, FOLIC ACID, CALCIUM PANTOTHENATE, BIOTIN, ASCORBIC ACID, CHOLENE CHLORIDE, INOSITOL, CALCIUM CARBOXYLATE, CALCIUM PHOSPHATE, MANGANESE SULFATE, COPPER SULFATE, POTASSIUM IODIDE, SODIUM SELENITE, SODIUM CHLORIDE, POTASSIUM CITRATE, POTASSIUM HYDROXIDE, TAURINE, L-CARNITINE, NUCLEOTIDES: URIDINE, 5'-MONOPHOSPHATE, CYTIDINE, 5'-MONOPHOSPHATE, DISODIUM GUANOSINE, 5'-MONOPHOSPHATE, DISODIUM URIDINE, 5'-MONOPHOSPHATE.

MEAD JOHNSON & COMPANY
EVANSVILLE, IN 47711 U.S.A.
©2009 Mead Johnson & Company

*A SOURCE OF ARACHIDONIC ACID (ARA)
**A SOURCE OF DOCOSAHENAIC ACID (DHA)

