


Adult Pneumococcal Vaccination Table



The Advisory Committee on Immunization Practices (ACIP) recommends all adults 65 years of age and older receive both PCV13 and PPSV23 pneumococcal vaccinations. ACIP also recommends adults 19 through 64 years of age who have underlying medical conditions receive pneumococcal vaccination with PCV13, PPSV23, or both. A second dose of PPSV23 before age 65 is not recommended for adults with Cerebral Spinal Fluid Leaks or adults with cochlear implants.

RISK GROUP	UNDERLYING MEDICAL CONDITION	PCV13	PPSV23	
		RECOMMENDED	RECOMMENDED	REVACCINATION AT 5 YEARS AFTER FIRST DOSE
Immunocompetent persons	Chronic heart disease: <i>congestive heart failure and cardiomyopathies</i>		✓	
	Chronic lung disease: <i>chronic obstructive lung disease, emphysema, and asthma</i>		✓	
	Diabetes mellitus		✓	
	Cerebral Spinal Fluid Leaks	✓	✓	
	Cochlear implants	✓	✓	
	Alcoholism		✓	
	Chronic liver disease		✓	
	Cigarette smoking		✓	
Persons with functional or anatomic asplenia	Sickle cell disease/other hemoglobinopathies	✓	✓	✓
	Congenital or acquired asplenia	✓	✓	✓
Immunocompromised persons	Congenital or acquired immunodeficiencies: <i>includes B-humoral or T-lymphocyte deficiency; complement deficiencies sC1, C2, C3 and C4 deficiencies; and phagocytic disorders, excluding chronic granulomatous disease</i>	✓	✓	✓
	HIV infection	✓	✓	✓
	Chronic renal failure	✓	✓	✓
	Nephrotic syndrome	✓	✓	✓
	Leukemia	✓	✓	✓
	Lymphoma	✓	✓	✓
	Hodgkin disease	✓	✓	✓
	Generalized malignancy	✓	✓	✓
	Latrogenic immunosuppression: <i>diseases requiring treatment with immunosuppressive drugs, long-term systemic corticosteroids, and radiation therapy</i>	✓	✓	✓
	Solid organ transplant	✓	✓	✓
	Multiple myeloma	✓	✓	✓




Adult Pneumococcal Vaccination Table

The Advisory Committee on Immunization Practices (ACIP) recommends all adults 65 years of age and older receive both PCV13 and PPSV23 pneumococcal vaccinations. ACIP also recommends adults 19 through 64 years of age who have underlying medical conditions receive pneumococcal vaccination with PCV13, PPSV23, or both. A second dose of PPSV23 before age 65 is not recommended for adults with Cerebral Spinal Fluid Leaks or adults with cochlear implants.



Source: Centers for Disease Control and Prevention. For more information visit www.cdc.gov/vaccines/vpd-vac/pneumo/vac-PCV13-adults.htm.



You never outgrow the need for vaccines.