





At this SOC or ROC	
Anterior	Posterior
	

(M1311) Current Number of Unhealed Pressure Ulcers at Each Stage	Enter Number
A1. Stage 2: Partial thickness loss of dermis presenting as a shallow open ulcer with red pink wound bed, without slough. May also present as an intact or open/ruptured blister. Number of Stage 2 pressure ulcers [If 0 at FU/DC Go to M1311B1]	<input type="text"/>
B1. Stage 3: Full thickness tissue loss. Subcutaneous fat may be visible but bone, tendon, or muscle is not exposed. Slough may be present but does not obscure the depth of tissue loss. May include undermining and tunneling. Number of Stage 3 pressure ulcers [If 0 at FU/DC Go to M1311C1]	<input type="text"/>
C1. Stage 4: Full thickness tissue loss with exposed bone, tendon, or muscle. Slough or eschar may be present on some parts of the wound bed. Often includes undermining and tunneling. Number of Stage 4 pressure ulcers [If 0 at FU/DC Go to M1311D1]	<input type="text"/>
D1. Unstageable: Non-removable dressing: Known but not stageable due to non-removable dressing/device Number of unstageable pressure ulcers due to non-removable dressing/device [If 0 at FU/DC Go to M1311E1]	<input type="text"/>
E1. Unstageable: Slough and/or eschar: Known but not stageable due to coverage of wound bed by slough and/or eschar Number of unstageable pressure ulcers due to coverage of wound bed by slough and/or eschar [If 0 at FU/DC Go to M1311F1]	<input type="text"/>
F1. Unstageable: Deep tissue injury: Suspected deep tissue injury in evolution Number of unstageable pressure ulcers with suspected deep tissue injury in evolution [If 0 - Go to M1322 (at Follow up), Go to M1313 (at Discharge)]	<input type="text"/>