

Missouri Resident Diabetes Profile

Indicator	Rate Denominator & Constant	Rate Type
Risk Factor for diabetes		
Among adults without diabetes (County Level Study)		
<i>Overweight without diabetes</i>	per 100 population	weighted percent
<i>Obesity without diabetes</i>	per 100 population	weighted percent
<i>Physical inactivity without diabetes</i>	per 100 population	weighted percent
Among youth (YTS/YRBSS)		
<i>Insufficient physical activity- High School</i>	per 100 population	weighted percent
<i>Insufficient physical activity- Middle School</i>	per 100 population	weighted percent
<i>Low fruit and vegetable intake- High School</i>	per 100 population	weighted percent
<i>Low fruit and vegetable intake- Middle School</i>	per 100 population	weighted percent
Diabetes prevalence among adults (BRFSS)		
<i>All adults >= 18 years of age</i>	per 100 population	weighted percent
<i>Age 18-44</i>	per 100 population	weighted percent
<i>Age 45-54</i>	per 100 population	weighted percent
<i>Age 55-64</i>	per 100 population	weighted percent
<i>Age 65+</i>	per 100 population	weighted percent
Annual household income		
<i>Less than \$15000</i>	per 100 population	weighted percent
<i>\$15000-\$24999</i>	per 100 population	weighted percent
<i>\$25000-\$34999</i>	per 100 population	weighted percent
<i>\$35000-\$49999</i>	per 100 population	weighted percent
<i>\$50000+</i>	per 100 population	weighted percent
Educational attainment		
<i>Less than high school</i>	per 100 population	weighted percent
<i>High school or GED</i>	per 100 population	weighted percent
<i>Some post high school</i>	per 100 population	weighted percent
<i>College or tech school graduate</i>	per 100 population	weighted percent
Risk factor complications among adults with diabetes (CLS)		
<i>Overweight with diabetes</i>	per 100 population	weighted percent
<i>Obesity with diabetes</i>	per 100 population	weighted percent
<i>Current smoking with diabetes</i>	per 100 population	weighted percent
<i>Physical inactivity with diabetes</i>	per 100 population	weighted percent

Missouri Resident Diabetes Profile

Indicator	Rate Denominator & Constant	Rate Type
<i>Low fruit and vegetable intake with diabetes</i>	per 100 population	weighted percent
Preventive care practices among adults with diabetes (BRFSS)		
<i>Teeth cleaned within past year</i>	per 100 population	weighted percent
<i>Current health coverage</i>	per 100 population	weighted percent
<i>A1C tests within the past year- 0 to 1 tests</i>	per 100 population	weighted percent
<i>A1C tests within the past year- 2 to 3 tests</i>	per 100 population	weighted percent
<i>A1C tests within the past year- 4 or more tests</i>	per 100 population	weighted percent
<i>Annual foot exam by a health care professional</i>	per 100 population	weighted percent
<i>Last dilated eye exam- within the last year</i>	per 100 population	weighted percent
<i>Last dilated eye exam- more than 1 year but less than 2 years</i>	per 100 population	weighted percent
<i>Last dilated eye exam- 2 or more years</i>	per 100 population	weighted percent
<i>Daily self checks for blood glucose</i>	per 100 population	weighted percent
<i>Flu shot within the past year</i>	per 100 population	weighted percent
<i>Pneumonia vaccination</i>	per 100 population	weighted percent
<i>Self-management education course</i>	per 100 population	weighted percent
Complications of diabetes for adults (BRFSS)		
<i>Eye complications and/or diabetic retinopathy</i>	per 100 population	weighted percent
Prev. care practices among adults with diabetes and hypertension (BRFSS)		
<i>Currently on high blood pressure medicine</i>	per 100 population	weighted percent
Mortality/PAS (Hospitalization and ER Indicators)		
Diabetes in pregnancy		
<i>Births to women with insulin-dependent diabetes</i>	per 1,000 resident live births	crude rate
<i>Births to women with other diabetes</i>	per 1,000 resident live births	crude rate
Emergency room visits for persons with diabetes		
<i>Diabetes as the principal diagnosis</i>	per year per 1,000 population	age-adjusted to the U.S. 2000 standard population
<i>Total charges</i>	-	-
<i>Diabetes as the principal or a secondary diagnosis</i>	per year per 1,000 population	age-adjusted to the U.S. 2000 standard population
Hospitalizations for persons with diabetes		
<i>Diabetes as the principal diagnosis</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Total charges</i>	-	-
<i>Days of care</i>	-	-

Missouri Resident Diabetes Profile

Indicator	Rate Denominator & Constant	Rate Type
<i>Diabetes as the principal or a secondary diagnosis</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Total charges</i>	-	-
<i>Days of care</i>	-	-
<i>Heart disease as the principal diagnosis</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Ischemic heart disease as the principal diagnosis</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Lower extremity condition as the principal diagnosis</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Lower extremity amputation</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Acute myocardial infarction (AMI)</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Acute cerebrovascular disease</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
<i>Chronic renal failure</i>	per year per 10,000 population	age-adjusted to the U.S. 2000 standard population
Mortality		
<i>Diabetes as underlying cause</i>	per year per 100,000 population	age-adjusted to the U.S. 2000 standard population
<i>Diabetes as underlying or contributing cause</i>	per year per 100,000 population	age-adjusted to the U.S. 2000 standard population