

2006 Summary of HIV Disease in Missouri

Figure 1. Persons living with HIV disease, newly diagnosed with HIV disease, AIDS related deaths, Missouri, 1997-2006



| Table 1. Percent of annual living HIV disease cases newly diagnosed in each year, Missouri, 1997-2006 | | | | | | | | | | | | |
|---|-------|------|------|------|------|------|------|------|------|------|--|--|
| | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | | |
| New Cases Percentage | 10.7% | 9.0% | 7.3% | 6.7% | 6.9% | 7.0% | 5.9% | 6.0% | 5.3% | 5.1% | | |

Note: Indicates prevention efforts have reduced the rate HIV transmission from persons living with HIV to those persons not infected with HIV.



^{**}Cases are indicated by year of initial diagnosis reported to the Missouri Department of Health and Senior Services. (The year in which the first diagnosis of the person, whether as an HIV case or an AIDS case, was documented by the Department.)

***These cases were either: 1) initially reported as HIV cases and then later reclassified as AIDS cases because they subsequently met the AIDS

case definition; or 2) initially reported as AIDS cases.





| | | HIV | | A STATE OF THE STA | AIDS | Н | IV Diseas | e** | |
|---------------------------------|-------|---------|------|--|---------|------|-----------|---------|------|
| Sex | Total | % | Rate | Total | % | Rate | Total | % | Rate |
| Male | 328 | 82.4% | 11.7 | 102 | 83.6% | 3.6 | 430 | 82.7% | 15.3 |
| Female | 70 | 17.6% | 2.4 | 20 | 16.4% | 0.7 | 90 | 17.3% | 3.1 |
| Total | 398 | 100.0% | 6.9 | 122 | 100.0% | 2.1 | 520 | 100.0% | 9.0 |
| Total | 000 | 100.070 | 0.0 | | 100.070 | | 020 | 100.070 | 0.0 |
| Age Group | Total | % | Rate | Total | % | Rate | Total | % | Rate |
| 0-1 | 3 | 0.8% | 2.0 | 0 | 0.0% | 0.0 | 3 | 0.6% | 2.0 |
| 2-12 | 2 | 0.5% | 0.2 | 0 | 0.0% | 0.0 | 2 | 0.4% | 0.2 |
| 13-18 | 19 | 4.8% | 3.8 | 1 | 0.8% | 0.2 | 20 | 3.8% | 4.0 |
| 19-24 | 71 | 17.8% | 13.7 | 8 | 6.6% | 1.5 | 79 | 15.2% | 15.2 |
| 25-44 | 227 | 57.0% | 14.4 | 76 | 62.3% | 4.8 | 303 | 58.3% | 19.2 |
| 45-64 | 72 | 18.1% | 5.1 | 33 | 27.0% | 2.3 | 105 | 20.2% | 7.4 |
| 65+ | 4 | 1.0% | 0.5 | 4 | 3.3% | 0.5 | 8 | 1.5% | 1.0 |
| Total | 398 | 100.0% | 6.9 | 122 | 100.0% | 2.1 | 520 | 100.0% | 9.0 |
| Race/Ethnicity and Sex | Total | % | Rate | Total | % | Rate | Total | % | Rate |
| White Male | 174 | 43.7% | 7.4 | 43 | 35.2% | 1.8 | 217 | 41.7% | 9.2 |
| Black Male | 137 | 34.4% | 43.3 | 43 | 35.2% | 13.6 | 180 | 34.6% | 56.9 |
| Hispanic Male | 8 | 2.0% | 10.2 | 15 | 12.3% | 19.1 | 23 | 4.4% | 29.3 |
| Asian Male | 1 | 0.3% | 2.5 | 1 | 0.8% | 2.5 | 2 | 0.4% | 5.0 |
| American Indian Male | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 |
| Unknown Race Male | 8 | 2.0% | N/A | 0 | 0.0% | N/A | 8 | 1.5% | N/A |
| White Female | 23 | 5.8% | 0.9 | 4 | 3.3% | 0.2 | 27 | 5.2% | 1.1 |
| Black Female | 41 | 10.3% | 11.5 | 15 | 12.3% | 4.2 | 56 | 10.8% | 15.7 |
| Hispanic Female | 3 | 0.8% | 4.3 | 1 | 0.8% | 1.4 | 4 | 0.8% | 5.7 |
| Asian Female | 2 | 0.5% | 4.7 | 0 | 0.0% | 0.0 | 2 | 0.4% | 4.7 |
| American Indian Female | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 |
| Unknown Race Female | 1 | 0.3% | N/A | 0 | 0.0% | N/A | 1 | 0.2% | N/A |
| Total | 398 | 100.0% | 6.9 | 122 | 100.0% | 2.1 | 520 | 100.0% | 9.0 |
| Exposure Category | Total | % | Rate | Total | % | Rate | Total | % | Rate |
| Men who have sex with men (MSM) | 195 | 49.0% | N/A | 62 | 50.8% | N/A | 257 | 49.4% | N/A |
| MSM/IDU | 11 | 2.8% | N/A | 3 | 2.5% | N/A | 14 | 2.7% | N/A |
| Injection drug use (IDU) | 18 | 4.5% | N/A | 6 | 4.9% | N/A | 24 | 4.6% | N/A |
| Heterosexual contact | 29 | 7.3% | N/A | 10 | 8.2% | N/A | 39 | 7.5% | N/A |
| Other risk | 0 | 0.0% | N/A | 1 | 0.8% | N/A | 1 | 0.2% | N/A |
| Unknown/Unreported risk | 140 | 35.2% | N/A | 40 | 32.8% | N/A | 180 | 34.6% | N/A |
| Pediatric risk | 5 | 1.3% | N/A | 0 | 0.0% | N/A | 5 | 1.0% | N/A |
| Total | 398 | 100.0% | 6.9 | 122 | 100.0% | 2.1 | 520 | 100.0% | 9.0 |
| HIV Region | Total | % | Rate | Total | % | Rate | Total | % | Rat |
| St Louis | 178 | 44.7% | 8.7 | 51 | 41.8% | 2.5 | 229 | 44.0% | 11. |
| Kansas City | 139 | 34.9% | 11.6 | 43 | 35.2% | 3.6 | 182 | 35.0% | 15.2 |
| Northwest | 6 | 1.5% | 2.5 | 1 | 0.8% | 0.4 | 7 | 1.3% | 2.9 |
| North Central | 15 | 3.8% | 2.1 | 9 | 7.4% | 1.2 | 24 | 4.6% | 3.3 |
| Southwest | 32 | 8.0% | 3.0 | 16 | 13.1% | 1.5 | 48 | 9.2% | 4.6 |
| Southeast | 4 | 1.0% | 0.8 | 10 | 0.8% | 0.2 | 5 | 1.0% | 1.0 |
| Corrections | 24 | 6.0% | N/A | 1 | 0.8% | N/A | 25 | 4.8% | N/A |
| Total | 398 | 100.0% | 6.9 | 122 | 100.0% | 2.1 | 520 | 100.0% | 9.0 |

^{*}Persons residing in Missouri at time of diagnosis.

**The sum of newly diagnosed HIV cases and newly diagnosed AIDS cases.

Note: Percentages may not total due to rounding.

Rates are per 100,000 residents, based on 2004 MDHSS estimates.

| Та | ble 3. | 3. Persons living with HIV disease, Missouri*, 2006 | | | | | | | | | |
|-------------------------------|--------|---|--------|-------------|-------|--------|--|------------------|-----------|--------------|--|
| | | | HIV | | | AIDS | and the second s | HI | / Disease | ** | |
| Sex | | Total | % | Rate | Total | % | Rate | Total | % | Rate | |
| Male | | 3,935 | 81.5% | 140.0 | 4,542 | 85.7% | 161.6 | 8,477 | 83.7% | 301.6 | |
| Female | | 895 | 18.5% | 30.4 | 759 | 14.3% | 25.8 | 1,654 | 16.3% | 56.2 | |
| Total | | 4,830 | 100.0% | 83.9 | 5,301 | 100.0% | 92.1 | 10,131 | 100.0% | 176.0 | |
| Age Group | | Total | % | Rate | Total | % | Rate | Total | % | Rate | |
| 0-1 | | 40 | 0.8% | 26.4 | 12 | 0.2% | 7.9 | 52 | 0.5% | 34.3 | |
| 2-12 | | 14 | 0.3% | 1.7 | 9 | 0.2% | 1.1 | 23 | 0.2% | 2.8 | |
| 13-18 | | 158 | 3.4% | 31.9 | 41 | 0.8% | 8.3 | 199 | 2.1% | 40.2 | |
| 19-24 | | 823 | 17.5% | 158.3 | 368 | 7.4% | 70.8 | 1,191 | 12.3% | 229.1 | |
| 25-44 | | 3,233 | 68.6% | 204.5 | 3,921 | 79.0% | 248.0 | 7,154 | 73.9% | 452.4 | |
| 45-64 | | 544 | 11.5% | 38.3 | 914 | 18.4% | 64.4 | 1,458 | 15.1% | 102.8 | |
| 65+ | | 18 | 0.4% | 2.4 | 36 | 0.7% | 4.7 | 54 | 0.6% | 7.1 | |
| Total | | 4,830 | 102.5% | 83.9 | 5,301 | 106.8% | 92.1 | 10,131 | 104.7% | 176.0 | |
| Race/Ethnicity and Sex | | Total | % | Rate | Total | % | Rate | Total | % | Rate | |
| White Male | | 2,164 | 44.8% | 91.7 | 2,653 | 50.0% | 112.4 | 4,817 | 47.5% | 204.0 | |
| Black Male | | 1,595 | 33.0% | 504.1 | 1,729 | 32.6% | 546.5 | 3,324 | 32.8% | 1050.6 | |
| Hispanic Male | | 114 | 2.4% | 145.2 | 131 | 2.5% | 166.9 | 245 | 2.4% | 312.1 | |
| Asian Male | | 18 | 0.4% | 44.6 | 11 | 0.2% | 27.3 | WESTEROFFERENCES | 0.3% | 71.9 | |
| | | 7 | | | | | | 29 | | | |
| American Indian Male | | | 0.1% | 47.4 | 17 | 0.3% | 115.2 | 24 | 0.2% | 162.6 | |
| Unknown Race Male | | 37 | 0.8% | N/A | 1 | 0.0% | N/A | 38 | 0.4% | N/A | |
| White Female | | 332 | 6.9% | 13.5 | 251 | 4.7% | 10.2 | 583 | 5.8% | 23.7 | |
| Black Female | | 526 | 10.9% | 147.3 | 485 | 9.1% | 135.8 | 1,011 | 10.0% | 283.2 | |
| Hispanic Female | | 22 | 0.5% | 31.6 | 18 | 0.3% | 25.8 | 40 | 0.4% | 57.4 | |
| Asian Female | | 6 | 0.1% | 14.0 | 4 | 0.1% | 9.4 | 10 | 0.1% | 23.4 | |
| American Indian Female | | 0 | 0.0% | 0.0 | 1 | 0.0% | 6.9 | 1 | 0.0% | 6.9 | |
| Unknown Race Female | | 9 | 0.2% | N/A | 0 | 0.0% | N/A | 9 | 0.1% | N/A | |
| Total | | 4,830 | 100.0% | 83.9 | 5,301 | 100.0% | 92.1 | 10,131 | 100.0% | 176.0 | |
| Exposure Category | | Total | % | Rate | Total | % | Rate | Total | % | Rate | |
| Men who have sex with men (MS | SM) | 2,723 | 56.4% | N/A | 3,281 | 61.9% | N/A | 6,004 | 59.3% | N/A | |
| MSM/IDU | | 217 | 4.5% | N/A | 399 | 7.5% | N/A | 616 | 6.1% | N/A | |
| Injection drug use (IDU) | | 315 | 6.5% | N/A | 405 | 7.6% | N/A | 720 | 7.1% | N/A | |
| Heterosexual contact | | 780 | 16.1% | N/A | 748 | 14.1% | N/A | 1528 | 15.1% | N/A | |
| Other risk | | 34 | 0.7% | N/A | 67 | 1.3% | N/A | 101 | 1.0% | N/A | |
| Unknown/Unreported risk | | 707 | 14.6% | N/A | 374 | 7.1% | N/A | 1081 | 10.7% | N/A | |
| Pediatric risk | | 54 | 1.1% | N/A | 27 | 0.5% | N/A | 81 | 0.8% | N/A | |
| Total | | 4,830 | 100.0% | 83.9 | 5,301 | 100.0% | 92.1 | 10,131 | 100.0% | 176.0 | |
| HIV Region | | Total | % | Rate | Total | % | Rate | Total | % | Rate | |
| St Louis | | 2,320 | 49.2% | 112.9 | 2,598 | 52.3% | 126.4 | 4,918 | 50.8% | 239.3 | |
| Kansas City | | 1,447 | 30.7% | 121.0 | 1,633 | 32.9% | 136.5 | 3,080 | 31.8% | 257.5 | |
| Northwest | | 42 | 0.9% | 17.4 | 73 | 1.5% | 30.3 | 115 | 1.2% | 47.7 | |
| North Central | | 189 | 4.0% | 26.2 | 243 | 4.9% | 33.6 | 432 | 4.5% | 59.8 | |
| Southwest | | 351 | 7.4% | 33.3 | 373 | 7.5% | 35.4 | 724 | 7.5% | 68.7 | |
| Southeast | | 125 | 2.7% | 25.7 | 147 | 3.0% | 30.2 | 272 | 2.8% | 56.0 | |
| Corrections | | 356 | 7.6% | 25.7 N/A | 234 | 4.7% | N/A | 590 | 6.1% | | |
| Total | | 4,830 | 102.5% | 83.9 | 5,301 | 106.8% | 92.1 | 10,131 | 104.7% | N/A 176.0 | |

^{*}Persons residing in Missouri at time of diagnosis.

**The sum of HIV cases and AIDS cases.

Note: Percentages may not total due to rounding.

Rates are per 100,000 residents, based on 2004 MDHSS estimates.

| | All New HIV+ Diagnosed in 2006 | | All New HIV+ Diagnosed in 2006 who had a CD4/VL | | All New HIV+ Diagnosed in 2006 with an Initial Diagnosis of AIDS | | All New HIV+ Diagnosed in 2006 (Case Management) | | All New HIV+ Diagnosed in 2006 who had a CD4/VL (Case Management) | | All New HIV+ Diagnosed in 2006 with an Initial Diagnosis of AIDS (Case Management | |
|-----------------|-----------------------------------|------------|---|------------|---|------------|--|------------|--|--------|---|--------|
| | # | % of Total | # | % of Total | # | % of Total | # | % of Total | # | %in CM | # | %in al |
| Total | 520 | 100.0% | 330 | 63.5% | 122 | 23.5% | 218 | 41.9% | 162 | 74.3% | 61 | 28.0% |
| Sex | | | | | | | | | | | | |
| Male | 430 | 100.0% | 274 | 63.7% | 102 | 23.7% | 173 | 40.2% | 129 | 74.6% | 50 | 28.9% |
| Female | 90 | 100.0% | 56 | 62.2% | 20 | 22.2% | 45 | 50.0% | 33 | 73.3% | 11 | 24.4% |
| Race/Ethnicity | | | | | | | | | | | | |
| White | 244 | 100.0% | 170 | 69.7% | 47 | 19.3% | 99 | 40.6% | 76 | 76.8% | 20 | 20.2% |
| Black | 236 | 100.0% | 130 | 55.1% | 58 | 24.6% | 106 | 44.9% | 75 | 70.8% | 34 | 32.1% |
| Hispanic | 27 | 100.0% | 22 | 81.5% | 16 | 59.3% | 12 | 44.4% | 10 | 83.3% | 7 | 58.3% |
| American Indian | 0 | - | 0 | - | 0 | _ | 0 | 1 | 0 | - | 0 | - |
| Asian | 4 | 100.0% | 3 | 75.0% | 1 | 25.0% | 0 | 0.0% | 0 | - | 0 | |
| Other/Unknown | 9 | 100.0% | 5 | 55.6% | 0 | 0.0% | 1 | 11.1% | 1 | 100.0% | 0 | 0.0% |
| Age Group | | | | | | | | | | | | |
| 0-1 | 3 | 100.0% | 2 | 66.7% | 0 | 0.0% | 0 | 0.0% | 0 | | 0 | - |
| 2 - 12 | 2 | 100.0% | 2 | 100.0% | 0 | 0.0% | 0 | 0.0% | 0 | - | 0 | - |
| 13 - 18 | 20 | 100.0% | 12 | 60.0% | 1 | 5.0% | 10 | 50.0% | 9 | 90.0% | 0 | 0.0% |
| 19 - 24 | 79 | 100.0% | 38 | 48.1% | 8 | 10.1% | 42 | 53.2% | 24 | 57.1% | 6 | 14.3% |
| 25 - 44 | 303 | 100.0% | 200 | 66.0% | 76 | 25.1% | 127 | 41.9% | 99 | 78.0% | 39 | 30.7% |
| 45 - 64 | 105 | 100.0% | 71 | 67.6% | 33 | 31.4% | 39 | 37.1% | 30 | 76.9% | 16 | 41.0% |
| 65+ | 8 | 100.0% | 5 | 62.5% | 4 | 50.0% | 0 | 0.0% | 0 | | 0 | - |
| Region | | | | | | | | | | | | |
| Kansas City | 182 | 100.0% | 134 | 73.6% | 43 | 23.6% | 75 | 41.2% | 68 | 90.7% | 24 | 32.0% |
| Northwest | 7 | 100.0% | 5 | 71.4% | 1 | 14.3% | 4 | 57.1% | 3 | 75.0% | 1 | 25.0% |
| North Central | 24 | 100.0% | 22 | 91.7% | 9 | 37.5% | 7 | 29.2% | 7 | 100.0% | 1 | 14.3% |
| St. Louis | 229 | 100.0% | 106 | 46.3% | 51 | 22.3% | 93 | 40.6% | 49 | 52.7% | 26 | 28.0% |
| Southwest | 48 | 100.0% | 43 | 89.6% | 16 | 33.3% | 29 | 60.4% | 29 | 100.0% | 8 | 27.6% |
| Southeast | 5 | 100.0% | 3 | 60.0% | 1 | 20.0% | 4 | 80.0% | 2 | 50.0% | 1 | 25.0% |
| Corrections | 25 | 100.0% | 17 | 68.0% | 1 | 4.0% | 6 | 24.0% | 4 | 66.7% | 0 | 0.0% |