

Health Alert:**Health Alert
April 20, 2018****Possible Measles
Exposures in the
Kansas City Area****FROM: RANDALL W. WILLIAMS, MD, FACOG
DIRECTOR****SUBJECT: Possible Measles Exposures in the Kansas City Area****April 20, 2018**

This document will be updated as new information becomes available. The current version can always be viewed at <http://www.health.mo.gov>.

The Missouri Department of Health & Senior Services (DHSS) is now using 4 types of documents to provide important information to medical and public health professionals, and to other interested persons:

Health Alerts convey information of the highest level of importance which warrants immediate action or attention from Missouri health providers, emergency responders, public health agencies, and/or the public.

Health Advisories provide important information for a specific incident or situation, including that impacting neighboring states; may not require immediate action.

Health Guidances contain comprehensive information pertaining to a particular disease or condition, and include recommendations, guidelines, etc. endorsed by DHSS.

Health Updates provide new or updated information on an incident or situation; can also provide information to update a previously sent Health Alert, Health Advisory, or Health Guidance; unlikely to require immediate action.

The Missouri Department of Health and Senior Services (DHSS) continues to receive reports of measles cases in Kansas City, Missouri. Newly identified exposures of concern include:

Date	Location Name	Exposure Timeframe	Location Address
April 6, 2018	KCI Expo Center-Midwest Parent Educators Vendor Hall	8:30 AM–1:00 PM	11730 NW Ambassador Dr. Kansas City, MO
April 10, 2018	Nebraska Furniture Mart	9:30 AM–2:00 PM	1601 Village West Pkwy Kansas City, KS

Note: Locations where individuals may have been exposed to measles, but can be identified, are not listed. Those individuals will be notified separately.

The Kansas City, Missouri Health Department and DHSS continue to work to identify and contact persons known to be potentially exposed to measles. However, potential transmission of the measles virus to unknown susceptible persons who had contact with the case may have occurred.

Measles is a highly contagious, acute viral illness that is transmitted by contact with an infected person through coughing and sneezing. Patients are considered to be contagious from 4 days before until 4 days after the rash appears. **Immune Globulin (IG)** can be administered to exposed individuals within 6 days of exposure to prevent or reduce the symptoms of measles infection for those without evidence of immunity. Health care providers that may see patients concerned about an exposure should consider having IG on hand in the event it is needed, both for this contact investigation event and in the future.

Health care providers should maintain a high index of suspicion for measles among febrile patients with a rash. Patients with clinical signs/symptoms compatible with measles (febrile rash plus cough, coryza, and/or conjunctivitis) should be asked about the exposures detailed above, recent travel abroad and contact with returning travelers, or contact with someone with a febrile rash illness. Their vaccination status should also be verified. Immunocompromised patients may not exhibit a rash, or may exhibit an atypical rash. The incubation period for measles from exposure to fever is usually about 10-12 days and from exposure to rash onset is usually 14 days (range, 7 to 21 days).

Office of the Director
912 Wildwood
P.O. Box 570
Jefferson City, MO 65102
Telephone: 800-392-0272
Fax: 573-751-6041

Website: <http://www.health.mo.gov>

Persons who have been exposed to measles should contact their health care provider if they develop cold-like symptoms with a fever and/or rash. They should **NOT** go to any health care facility without calling first. The suspect case should be kept separated from others to prevent further spread. (Note that measles virus can remain infectious in the air for up to 2 hours after an infected person leaves an area such as a waiting room.) Isolate suspect measles case-patients and immediately report suspected cases to the local public health agency, or to DHSS at 573/751-6113 or 800/392-0272 (24/7). To ensure prompt public health response, do not wait for laboratory confirmation.

Detection of measles-specific IgM antibody and measles RNA by real-time polymerase chain reaction (RT-PCR) are the most common methods for confirming measles infection. Healthcare providers should obtain both a serum sample and a throat swab (or nasopharyngeal swab) from patients suspected to have measles at first contact with them.

The Missouri State Public Health Laboratory (MSPHL) provides laboratory support for the diagnosis of measles infections occurring in Missouri. MSPHL will only test specimens that are approved by state public health officials. A specimen for molecular detection by RT-PCR should be collected and submitted to MSPHL along with the initial serum specimen, and include NP swab, throat swab, or urine (see the CDC instructions below). Please note that a RT-PCR specimen should NOT be substituted for a serum specimen. Serum specimens submitted to MSPHL will be tested for measles IgM and rubella IgM as requested by the investigating epidemiologist. The RT-PCR specimen will be referred to a Vaccine Preventable Disease (VPD) Reference Laboratory. VPD laboratories are established in cooperation with public health laboratories and the Centers for Disease Control and Prevention [CDC] to provide reference testing and surge capacity.

The sensitivity of measles IgM assays varies, and may be diminished during the first 72 hours after rash onset. If the result is negative for measles IgM and the patient has a generalized rash lasting more than 72 hours, a second serum specimen should be obtained and the measles IgM test should be repeated (AAP. *Red Book*, 2015; p. 537).

For questions regarding storage and shipping of all samples, please contact the MSPHL Virology Unit at 573-751-3334.

Measles serology instructions:

<http://health.mo.gov/lab/measlesrubella.php>

CDC measles RT-PCR instructions (do NOT ship specimens directly to CDC):

<https://www.cdc.gov/measles/lab-tools/rt-pcr.html>

For further guidance, please refer to:

<https://www.cdc.gov/measles/hcp/index.html>

https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6204a1.htm?s_cid=rr6204a1_w

Questions should be directed to DHSS' Bureau of Communicable Disease Control and Prevention at 573/751-6113 or 800/392-0272 (24/7).