

LPHA's Reference Guide to Hepatitis Data Entry into WebSurv



<p style="text-align: center;">Acute Hepatitis B Suspect</p> <ul style="list-style-type: none"> HBsAg (+) IgM Anti-HBc (+) HBeAg (+) HBV DNA (+) 	<p style="text-align: center;">Chronic Hepatitis B Suspect</p> <ul style="list-style-type: none"> Anti-HBc Total (+) HBV DNA <357 Anti-HBe (+)
---	---

When a single laboratory result is received without any other hepatitis B test results available, please refer to the above list to determine the correct condition status to be entered into WebSurv.

<p style="text-align: center;">Acute Hepatitis B Confirmed</p> <ul style="list-style-type: none"> HBsAg (+) AND/OR IgM Anti-HBc (+) <u>AND</u> Discrete onset of symptoms (i.e. fatigue, adnominal pain, loss of appetite, nausea, vomiting, dark urine) <u>AND</u> Jaundice OR Elevated ALT/AST 	<p style="text-align: center;">Chronic Hepatitis B Confirmed</p> <p>Any combination of the following tests greater than six months apart:</p> <ul style="list-style-type: none"> HBsAg (+) HBeAg (+) HBV DNA (+)
---	--

Acute Uncomplicated

Chronic Uncomplicated



Perinatal Hepatitis B

- Complete the IMMP 29 (when (+) pregnancy status is first determined and return to the Bureau of HIV, STD and Hepatitis)
- Complete the IMMP 29A on every household and sexual contact to the positive hepatitis B mom and return to the Bureau of HIV, STD and Hepatitis
- Provide the Bureau HIV, STD and Hepatitis with an updated IMMP 29 when the infant receives additional hepatitis B vaccines

AND/OR

- Update the state's Immunization registry when the infant receives his/her hepatitis B vaccine

Please encourage the physician to order post-vaccination HBsAg and Anti-HBs serologic tests on every infant born to a woman infected with hepatitis B, 1-2 months after the infant(s) completes the hepatitis B vaccination series.

A single positive Anti-HCV test result is insufficient to assign a confirmed hepatitis C condition. Enter a single positive Anti-HCV test result into WebSurv as a chronic hepatitis C suspect. If additional hepatitis C tests results are received that meet the confirmatory case condition, change the condition status to the appropriate confirmed status according to the table below.

Chronic Hepatitis C Suspect

- Anti-HCV (+)

Chronic Hepatitis C Confirmed

- Anti-HCV with a S/CO ratio > the reference range
- RIBA (+)
- HCV RNA (+)
- HCV Genotype

Acute Hepatitis C Confirmed

- Anti-HCV (+)
AND
- Discreet onset of symptoms (i.e. anorexia, abdominal discomfort, nausea, vomiting.) Jaundice OR ALT >400 IU/L
AND
- IgM Anti-HAV (-)
And
- IgM Anti-HBV (-)