



# Missouri Weekly Influenza Surveillance Report 2016-2017 Influenza Season<sup>1</sup>

**Week 8: February 19 – February 25, 2017**

*All data are preliminary and may change as more reports are received.*

## Summary:

- The estimated influenza activity in Missouri is Widespread<sup>2</sup>.
- A season-to-date total of 44,961 laboratory-positive<sup>3</sup> influenza cases (34,712 influenza A, 9,362 influenza B, and 887 untyped) have been reported in Missouri as of Week 8. The influenza type for reported cases season-to-date includes 77% influenza A, 21% influenza B, and 2% untyped. The highest season-to-date rates of reported laboratory-positive influenza cases are among children aged 0-4 years (1,630 cases per 100,000 population) and 5-14 years (1,519 cases per 100,000). Eleven laboratory-confirmed case of influenza [eight influenza A (H3), two influenza B (Yamagata), and one influenza A (H1N1)] were reported by the Missouri State Public Health Laboratory (MSPHL) during Week 8.
- The Centers for Disease Control and Prevention (CDC) has antigenically characterized six influenza isolates from Missouri, to date, this influenza season. Four viruses were antigenically similar to the A/Hong Kong/4801/2014-like (H3N2) virus, one virus was antigenically similar to the B/Brisbane/60/2008-like virus, and one virus was antigenically similar to the B/Phuket/3073/2013-like virus. An A/Hong Kong/4801/2014-like (H3N2) virus and a B/Brisbane/60/2008-like virus are included in the 2016-2017 Northern Hemisphere trivalent and quadrivalent vaccine formulations. A B/Phuket/3073/2013-like virus is included in the 2016-2017 Northern Hemisphere quadrivalent vaccine formulation.
- Influenza-like illness (ILI) activity is above baseline for both the Missouri Outpatient ILI Surveillance Network (ILINet) and the hospital emergency room visit chief complaint data reported through ESSENCE. The reported percentage of visits for ILI was 4.82% and 4.25% through ILINet and ESSENCE respectively.<sup>4</sup> The percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased during Week 8.
- Thirty-seven influenza-associated deaths have been reported in Missouri as of Week 8. During Week 7, 87 deaths involving Pneumonia and Influenza (P&I) were reported to the Bureau of Vital Records, resulting in a season-to-date total of 1,301 P&I associated deaths in Missouri.<sup>5</sup>
- Thirty-eight influenza or ILI-associated outbreaks have been reported in Missouri as of Week 8. Eleven influenza or ILI-associated school closures have been reported in Missouri as of Week 8.
- Influenza activity decreased slightly but remained elevated in the U.S. during Week 7. National influenza surveillance information is prepared by CDC and is included in the weekly FluView report, which is available online at <http://www.cdc.gov/flu/weekly/fluactivitysurv.htm>.

<sup>1</sup>The 2016-2017 influenza season begins CDC Week 40 (week ending October 8, 2016) and ends CDC Week 39 (week ending September 30, 2017).

<sup>2</sup>Widespread is defined as: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

<sup>3</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

<sup>4</sup>Influenza-like illness (ILI) is defined by ILINet as fever (temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat without a known cause other than influenza. Influenza-like illness (ILI) is defined by ESSENCE as Emergency Department chief complaints that contain keywords such as “flu”, “flulike”, “influenza” or “fever” plus “cough” or “fever” plus “sore throat”.

<sup>5</sup>The P&I data are available one week later. The P&I data for the CDC Week provided is the most current data available.

## Surveillance Data:

### Interactive Maps

The county specific influenza data are provided through interactive maps available at <http://arcg.is/2lXDf0a>. Click on the county to view the influenza data specific to that county.

- Reported Laboratory-positive Influenza Cases by Influenza Type by County, CDC Week 8
- Reported Laboratory-positive Influenza Cases by Influenza Type by County, Season-to-Date
- Reported Rate per 100,000 Population, CDC Week 8

### Data Figures

**Figure 1. Number of Laboratory-positive<sup>†</sup> Influenza Cases by Influenza Type, Missouri, CDC Week 8 (February 19 – February 25, 2017)<sup>\*</sup>**

Influenza Type	Week 6	Week 7	Week 8	2016-2017* Season-to-Date
Influenza A	6,610	5,969	3,040	34,712
Influenza B	1,808	2,352	1,832	9,362
Influenza Unknown Or Untyped	196	132	53	887
<b>Total</b>	<b>8,614</b>	<b>8,453</b>	<b>4,925</b>	<b>44,961</b>

<sup>†</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*Influenza season begins week ending October 8, 2016 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv).

**Figure 2. Number of Laboratory-positive<sup>†</sup> Influenza Cases and Case Rates by Age Group, Missouri, CDC Week 8 (February 19 – February 25, 2017)<sup>\*\*</sup>**

Age Group	Week 8 Cases	Week 8 Rate <sup>‡</sup>	2016-2017* Season-to-Date	2016-2017* Season-to-Date Rate <sup>‡</sup>
00-04	769	205	6,101	1,630
05-14	1,472	188	11,871	1,519
15-64	1,983	50	20,740	522
65+	701	75	6,247	670
<b>Total</b>	<b>4,925</b>	<b>81</b>	<b>44,961</b>	<b>741</b>

<sup>†</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*Influenza season begins week ending October 8, 2016 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

<sup>‡</sup>Incidence Rate per 100,000 population

**Figure 3. Number of Laboratory-positive<sup>†</sup> Influenza Cases and Case Rates by Region, Missouri, CDC Week 8 (February 19 – February 25, 2017)<sup>\*\*</sup>**


District	Week 8 Cases	Week 8 Rate <sup>‡</sup>	2016-2017* Season-to-Date	2016-2017* Season-to-Date Rate <sup>‡</sup>
CE	405	61	4,731	714
EA	1,862	82	12,144	538
NW	1,543	97	16,000	1,005
SE	676	142	5,659	1,189
SW	439	41	6,427	597
<b>Total</b>	<b>4,925</b>	<b>81</b>	<b>44,961</b>	<b>741</b>

<sup>†</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*Influenza season begins week ending October 8, 2016 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

<sup>‡</sup>Incidence Rate per 100,000 population


**Figure 4. Number of Laboratory-positive<sup>†</sup> Influenza Cases by CDC Week, Missouri, 2013-2017<sup>\*</sup>**



<sup>†</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*2016-2017 season-to-date through the week ending May 20, 2017 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. Week 53, 2014-2015 flu season has 3,082 reported confirmed cases. Data Source: Missouri Health Information Surveillance System (WebSurv).

**Figure 5. Percentage of Outpatient Visits for Influenza-like-Illness (ILI), Missouri Outpatient ILI Surveillance Network (ILINet) 2013-2017\*†**




\*The ILINet Region 7 (MO, IA, KS, NE) baseline is the mean percentage of patient visits for ILI during non-influenza weeks for the previous three seasons, plus two standard deviations. A non-influenza week is defined as periods of two or more consecutive weeks in which each week accounted for less than 2% of the season's total number of specimens that tested positive for influenza.

Data Source: U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Centers for Disease Control and Prevention (CDC).


†2016-2017 season-to-date through the week ending May 20, 2017 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of outpatient visits for ILI during Week 53 was 7.63.

**Figure 6. Season-to-Date PCR (+) Tests for Influenza in Missouri**



Data Source: National Respiratory and Enteric Virus Surveillance System (NREVSS), Centers for Disease Control and Prevention (CDC). 2016-2017 season-to-date through the week ending May 20, 2017 (Week 20).


**Figure 7. Weekly Percentage of Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, for 2013-2017 Influenza Seasons\*†**



\*The ESSENCE ILI Baseline is the mean percent of ILI visits for each week during the previous three years (2013-15) when percentage of ILI visits were less than 2% of total visits, plus two standard deviations.  
 Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE.


†The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of visits for ILI in ESSENCE participating hospitals during Week 53 was 4.3.

**Figure 8. Percentage of Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by Age-group, District and Statewide, CDC Week 8, 2017**



Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE.

**Figure 9. Percentage of Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by District and Statewide, for the 2016-2017 Influenza Season<sup>\*†</sup>**




Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE.

\*Changes in ILI surveillance for the Northwest District were implemented on January 16, 2017.

† Not all data was available for the Northwest District during Week 6.

**Figure 10. Patients Hospitalized with Influenza and/or Pneumonia Syndromes from Participating Missouri Hospitals, by Age Group, CDC Week 8, 2017**



Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE. Population data from DHSS Population MICA 2014 (<http://health.mo.gov/data/mica/mica/population.php>).

**Additional Influenza Data Sources:**

**Centers for Disease Control and Prevention: National Influenza Surveillance (FluView):**  
<http://www.cdc.gov/flu/weekly/fluactivitysurv.htm>

**The National Respiratory and Enteric Virus Surveillance System (NREVSS):**  
<https://www.cdc.gov/surveillance/nrevss/>

**World Health Organization: International Influenza Surveillance:**  
[http://www.who.int/influenza/surveillance\\_monitoring/en/](http://www.who.int/influenza/surveillance_monitoring/en/)