

## **Emergency Use Instructions for Healthcare Providers: Pfizer-BioNTech COVID-19 vaccine for Primary, Additional, and/or Booster Doses**

The Centers for Disease Control and Prevention (CDC) is issuing Emergency Use Instructions (EUI) to provide information about the use of the COVID-19 vaccine by Pfizer-BioNTech (Comirnaty), which is approved (licensed) by the Food and Drug Administration (FDA) for the prevention of COVID-19 in individuals 16 years and older.<sup>1</sup> The CDC-issued EUI provide instructions and information for the use of this vaccine that are beyond the FDA-approved labeling. The CDC-issued EUI provide information on the following uses of the COVID-19 vaccine by Pfizer-BioNTech for:

- primary dose(s) in those with certain immunocompromising conditions or those with incomplete primary dose series, and/or a booster dose for persons 12 years and older who received primary vaccination with certain **non-FDA authorized or approved** COVID-19 vaccines<sup>2</sup>.
- a 3-month interval for the booster dose after an mRNA vaccine primary series for persons 12 years and older who are moderately or severely immunocompromised.
- an additional dose in persons 18 years and older with certain immunocompromising conditions who received primary vaccination with the Janssen COVID-19 Vaccine.
- revaccination of moderately or severely immunocompromised persons 12 years and older who received certain therapies (i.e., hematopoietic cell transplant (HCT) or chimeric antigen receptor (CAR)-T-cell therapy) and received dose(s) of COVID-19 vaccine prior to or during treatment.

mRNA vaccines are preferred for persons with moderate or severe immune compromise. The COVID-19 vaccine by Moderna under EUI also allow similar uses in persons 18 years and older as an alternative mRNA COVID-19 vaccine to Pfizer-BioNTech, and the same or similar recommendations in the EUI also apply to the use of the COVID-19 vaccine by Moderna under EUI. See the [Moderna EUI Fact Sheet for Healthcare Providers](#).

Refer to CDC's [Interim Clinical Considerations](#) for specific recommendations on use of the COVID-19 vaccine by Pfizer-BioNTech allowed under the EUI. Relevant information is detailed in the sections titled: "People who received COVID-19 vaccine outside the United States", "People who received COVID-19 vaccine as part of a clinical trial", and "Recommendations for COVID-19 vaccination in moderately or severely immunocompromised people." For additional information about the COVID-19 vaccine by Pfizer-BioNTech COVID-19, refer to the [Comirnaty](#) package insert or the [Full Emergency Use Authorization \(EUA\) Prescribing Information](#) (FDA, 2022).

### **What are EUI and why is CDC issuing EUI for the COVID-19 vaccine by Pfizer-BioNTech?**

In 2013, the Pandemic and All-Hazards Preparedness Reauthorization Act included a new provision that allowed for the issuance of EUI to permit CDC to inform healthcare providers and recipients about certain uses of FDA-approved or cleared medical products. Specifically, EUI inform healthcare providers and recipients about such products' approved, licensed, or cleared conditions of use. The CDC Director has statutory (legal) authority to create, issue, and disseminate EUI before or during an emergency.

The COVID-19 vaccine by Pfizer-BioNTech was approved by the FDA in August 2021 as a 2-dose primary series for active immunization to prevent COVID-19 in persons 16 years and older. CDC is issuing these EUI to provide information about use of the COVID-19 vaccine by Pfizer-BioNTech for persons 12 years and older as primary

---

<sup>1</sup> Comirnaty is the proprietary name for the product licensed under the Biologics License Application (BLA). The Pfizer-BioNTech COVID-19 Vaccine has been available since December 11, 2020, pursuant to Emergency Use Authorization (EUA). The two approved formulations of Comirnaty and the two FDA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine for ≥12 years are the same formulations, and vials of the BLA-compliant vaccine may bear the name "Pfizer-BioNTech COVID-19 Vaccine." Because of these features, and because Comirnaty is commonly referred to as the "Pfizer vaccine" or the "Pfizer-BioNTech COVID-19 Vaccine," these EUI refer to this vaccine as the COVID-19 vaccine by Pfizer-BioNTech.

<sup>2</sup> A non-FDA authorized or approved COVID-19 vaccine that is listed for emergency use by the World Health Organization, or is included in CDC's [Technical Instructions for Implementing Presidential Proclamation Advancing Safe Resumption of Global Travel During the COVID-19 Pandemic](#) and [CDC's Order](#), or that is a non-placebo part of a clinical trial within or outside the United States that is a WHO-EUL COVID-19 vaccine or a vaccine that is not listed for emergency use by WHO but for which a U.S. data and safety monitoring board or equivalent has independently confirmed efficacy in the United States (hereinafter "non-FDA authorized or approved COVID-19 vaccines").



dose(s) in those with certain immunocompromising conditions or those with incomplete primary dose series, and/or a booster dose in those who received primary vaccination with certain **non-FDA authorized or approved** COVID-19 vaccines. For example, these EUI cover use of the COVID-19 vaccine by Pfizer-BioNTech in individuals who were vaccinated outside of the United States or in clinical trials with the AstraZeneca COVID-19 vaccine, the Novavax COVID-19 vaccine, or the Sinopharm COVID-19 vaccine, among others. The EUI also provide for heterologous use of the COVID-19 vaccine by Pfizer-BioNTech as an additional dose in persons 18 years and older with certain immunocompromising conditions who completed primary vaccination with the FDA-authorized Janssen COVID-19 Vaccine. Additionally, the EUI provide for revaccination of moderately or severely immunocompromised persons 12 years and older who received certain therapies (i.e. HCT or CAR-T-cell therapy) and received dose(s) of COVID-19 vaccine prior to or during treatment.

### **What is COVID-19?**

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by the novel coronavirus, SARS-CoV-2, that emerged in late 2019. It is predominantly a respiratory illness that can affect other organs. People with SARS-CoV-2 infection have reported a wide range of symptoms, ranging from no symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Symptoms may include fever or chills, cough, shortness of breath, fatigue, muscle or body aches, headache, new loss of taste or smell, sore throat, congestion or runny nose, nausea or vomiting, and diarrhea.

### **Who can receive the COVID-19 vaccine by Pfizer-BioNTech?**

The below describes who can receive the COVID-19 vaccine by Pfizer-BioNTech under EUI. The COVID-19 vaccine by Moderna can also be used under EUI for similar uses in those 18 years and older as an alternative mRNA COVID-19 vaccine (see the [Moderna EUI Fact Sheet for Healthcare Providers](#)).

- Certain moderately and severely immunocompromised persons 12 years and older who received primary vaccination with certain **non-FDA authorized or approved** COVID-19 vaccines should receive an additional primary dose of the COVID-19 vaccine by Pfizer-BioNTech.
- Persons 12 years and older who received incomplete primary dose series (e.g., only the first dose of 2-dose primary series) with certain **non-FDA authorized or approved** COVID-19 vaccines should receive a primary dose of the COVID-19 vaccine by Pfizer-BioNTech.
- Persons 12 years and older who have received primary vaccination with certain **non-FDA authorized or approved** COVID-19 vaccines should receive a booster dose of the COVID-19 vaccine by Pfizer-BioNTech.
- Certain moderately and severely immunocompromised persons 18 years and older who received primary vaccination with the Janssen COVID-19 Vaccine should receive an additional dose with the COVID-19 vaccine by Pfizer-BioNTech.
- Moderately or severely immunocompromised persons 12 years and older who received certain therapies (i.e., HCT or CAR-T-cell therapy) and received dose(s) of COVID-19 vaccine prior to or during treatment should be revaccinated with the COVID-19 vaccine by Pfizer-BioNTech for any doses received before or during treatment.

## What are the doses and intervals of the COVID-19 vaccine by Pfizer-BioNTech for primary, additional, and/or booster doses?

- A primary dose, including as an additional primary dose for those with certain immunocompromising conditions, of the Pfizer-BioNTech vaccine (30 µg in 0.3 mL) should be administered intramuscularly at least 28 days after primary vaccination with certain non-FDA authorized or approved COVID-19 vaccines.
- A booster dose of the COVID-19 vaccine by Pfizer-BioNTech (30 µg in 0.3 mL) should be administered intramuscularly for persons who completed primary vaccination with a series that included certain non-FDA authorized or approved COVID-19 vaccines: at least 3 months after completion of primary vaccination for immunocompromised persons or at least 5 months after completion of primary vaccination for non-immunocompromised persons.
- An additional dose with the COVID-19 vaccine by Pfizer-BioNTech (30 µg in 0.3 mL) should be administered intramuscularly for persons with certain immunocompromising conditions at least 28 days after completion of primary vaccination with the Janssen COVID-19 Vaccine (e.g., 1 primary dose of the Janssen COVID-19 Vaccine followed by an additional dose with Pfizer-BioNTech vaccine at least 28 days after the primary dose). People who received 1 primary dose of the Janssen COVID-19 Vaccine and 1 booster of an authorized COVID-19 vaccine, should receive an additional dose with the COVID-19 vaccine by Pfizer-BioNTech at least 2 months after the booster dose.
- Revaccination with the COVID-19 vaccine by Pfizer-BioNTech at least 3 months after therapy for any doses received before or during treatment with certain therapies (i.e., HCT or CAR-T-cell therapy).

See [Table 3](#) COVID-19 vaccination schedule for moderately or severely immunocompromised people in [CDC's Interim Clinical Considerations](#) for the latest dosing recommendations.

On a case-by-case basis, providers of moderately or severely immunocompromised patients may administer the COVID-19 vaccine by Pfizer-BioNTech outside of the FDA-authorized or FDA-approved labeling and CDC recommended dosing intervals based on clinical judgment when the benefits of vaccination are deemed to outweigh the potential and unknown risks for the recipient.

Refer to CDC's [Interim Clinical Considerations](#) for specific and the latest dosing recommendations (e.g., number of doses, dosing intervals, revaccination) that may vary for individuals with certain medical conditions and/or in certain circumstances, which differ from or extend beyond the FDA-authorized and/or FDA-approved labeling.

## What are the formulations of the COVID-19 vaccine by Pfizer-BioNTech that these EUI apply to?

The EUI apply to the FDA-approved formulations of the COVID-19 vaccine by Pfizer-BioNTech. As of December 16, 2021, there are two FDA-approved formulations of this vaccine that are distinguished by purple and gray caps. They are also FDA-authorized under EUA. The multiple dose vials (MDVs) with purple caps contain phosphate buffered saline that must be diluted with normal saline prior to administration. The MDVs with gray caps contain tromethamine (Tris) buffer and do not require dilution for administration. Each formulation, when prepared according to its respective formulation-specific instructions for administration, provide 0.3 mL doses (each containing 30 µg mRNA). FDA has explained that these formulations of the vaccine can be used [interchangeably](#) without presenting any safety or effectiveness concerns, when prepared according to their respective formulation-specific instructions for use.

## What are the common side effects with the COVID-19 vaccine by Pfizer-BioNTech?

Adverse reactions following administration of the vaccine that have been reported in clinical trials and/or post authorization include injection site pain, fatigue, headache, muscle pain, chills, joint pain, fever, injection site swelling, injection site redness, nausea, malaise, lymphadenopathy, decreased appetite, rash, pain in extremity, diarrhea, and vomiting.

## What are possible serious side effects with the COVID-19 vaccine by Pfizer-BioNTech?

Severe allergic reactions, including anaphylaxis, and other hypersensitivity reactions (e.g., rash, pruritus, urticaria, angioedema), syncope, myocarditis and pericarditis have been reported following administration of the vaccine outside of clinical trials. Myocarditis and/or pericarditis are rare, serious adverse events that have been reported after receipt of mRNA COVID-19 vaccines, with the highest risk currently observed in males 12–29 years.

### **Who should not receive the COVID-19 vaccine by Pfizer-BioNTech?**

Do not administer the COVID-19 vaccine by Pfizer-BioNTech to persons with known history of a severe allergic reaction (e.g., anaphylaxis) to a previous dose or any component of the vaccine (see *Contraindications, and Warnings and Precautions* sections in the [Comirnaty](#) package insert or [Full EUA Prescribing Information](#) as well as CDC's [Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Approved or Authorized in the United States](#) for additional considerations).

### **What information should be provided to persons receiving a primary, additional, and/or booster dose of the COVID-19 vaccine by Pfizer-BioNTech as described in the EUI?**

- Provide the [EUI Fact Sheet for Recipients and Caregivers](#).
- Provide a CDC COVID-19 Vaccination Record Card to the recipient or their caregiver with the lot number and date of administration recorded for the primary, additional, or booster dose of the COVID-19 vaccine by Pfizer-BioNTech.
- Provide the v-safe information sheet to vaccine recipients/caregivers and encourage vaccine recipients to participate in v-safe. V-safe is a voluntary smartphone-based tool that uses text messaging and web surveys to check in with people who have been vaccinated to identify potential side effects after COVID-19 vaccination. V-safe asks questions that help CDC monitor the safety of COVID-19 vaccines. For more information, visit: [www.cdc.gov/vsafe](http://www.cdc.gov/vsafe).

### **What is the available supporting evidence for use of the COVID-19 vaccine by Pfizer-BioNTech for additional primary or booster doses in people who received a primary vaccination with non-FDA authorized or approved COVID-19 vaccines?**

CDC has not systematically evaluated the safety, immunogenicity, and efficacy of an additional dose of the COVID-19 vaccine by Pfizer-BioNTech (as either an additional primary dose for immunocompromised or as a booster dose) following receipt of primary vaccination with a non-FDA authorized or approved COVID-19 vaccine. However, studies of COVID-19 vaccine boosting in the United Kingdom have shown that a third dose of AstraZeneca, Moderna, or Pfizer-BioNTech COVID-19 vaccines successfully boosted immune responses in people who had been primed with two doses of Pfizer-BioNTech or AstraZeneca COVID-19 vaccines approximately 3 months earlier. Levels of binding (IgG) and neutralizing antibodies, including against Delta variant, were generally higher when an mRNA vaccine was used as either a heterologous or homologous boost, or where the AstraZeneca COVID-19 vaccine was used as a heterologous boost after primary vaccination with the Pfizer-BioNTech COVID-19 vaccine (Munro et al., 2021). Frequencies of local and systemic adverse reactions in the 7 days post booster vaccination were higher with heterologous than homologous boosters and in those aged under 70 years when compared to older recipients. Frequencies of local and systemic adverse reactions were higher when the AstraZeneca COVID-19 vaccine was used to boost those who received primary vaccination with the Pfizer-BioNTech COVID-19 vaccine, when compared with the Pfizer-BioNTech COVID-19 vaccine after either primary vaccination (Munro et al., 2021).

Additional supporting evidence for use of the COVID-19 vaccine by Pfizer-BioNTech for additional primary or booster doses in people who received a non-FDA authorized or approved COVID-19 primary vaccination series are as follows. An unpublished small, randomized trial in Bahrain found that a third dose of Pfizer-BioNTech COVID-19 vaccine after a 2-dose Sinopharm BIBP COVID-19 vaccine primary series resulted in higher levels of IgG antibodies against the spike-antigen of SARS-CoV-2 (anti-S-IgG) compared to a 3-dose series of Sinopharm

BIBP COVID-19 vaccine (SAGE, 2021). In a pilot prospective cohort study of healthcare workers (HCWs) from Lebanon, 50 HCWs who received a 2-dose primary series of Sinopharm BIBP COVID-19 vaccine and a single booster dose of Pfizer-BioNTech COVID-19 vaccine had significantly higher anti-S-IgG titers compared to 50 homologous vaccinees (2 primary series doses and 1 booster dose of Pfizer-BioNTech COVID-19 vaccine) (Moghnieh, 2021). A longitudinal study of 41 Thai HCWs who received a 2-dose primary series of Sinovac (CoronaVac) COVID-19 vaccine demonstrated booster antibody responses following either AstraZeneca or Pfizer-BioNTech COVID-19 vaccines, including against the Delta variant (Patamatamkul, 2021). Local and systemic reactogenicity was reported to be mild to moderate across studies. Finally, a study from Chile examining heterologous boosting with AstraZeneca or Pfizer-BioNTech among Sinovac-CoronaVac primed individuals demonstrated higher vaccine effectiveness against infection, symptomatic disease, and intensive care unit admission compared with homologous boosting (Araos, 2021).

WHO's Strategic Advisory Group of Experts (SAGE) on Immunization has noted that although data are currently limited on the safety, immunogenicity, and effectiveness of heterologous versus homologous additional doses, evolving evidence suggests that use of a heterologous vaccine for an additional dose may be more immunogenic than a homologous series. In its recommendations for an additional dose in certain immunocompromised people and in people aged 60 years and over who received Sinopharm BIBP or Sinovac-CoronaVac COVID-19 vaccines as a 2-dose primary series, WHO has advised that countries can consider heterologous additional doses based on supply availability (WHO SAGE 2021a-c).

More than 80 countries are using boosters after non-FDA authorized or approved COVID-19 vaccines. Countries such as the United Kingdom (JCVI, 2021a-b), Canada (National Advisory Committee on Vaccination, 2021), Germany, and France have recommended heterologous dosing, including with use of Pfizer-BioNTech COVID-19 vaccine, for an additional primary series and/or booster dose based on their reviews of available immunological and safety data, as well as the epidemiology of COVID-19 and other contextual factors.

The heterologous booster dose of Pfizer-BioNTech COVID-19 vaccine in individuals who completed primary vaccination with Janssen COVID-19 Vaccine is supported by the immunogenicity data from a Phase 1/2 open-label clinical study ([NCT04889209](https://clinicaltrials.gov/ct2/show/study/NCT04889209)) by the National Institutes of Health conducted in the United States that evaluated heterologous booster regimens of FDA-authorized COVID-19 vaccines. In this study, adults who had completed primary vaccination with a Moderna COVID-19 Vaccine 2-dose series (N=151), a Janssen COVID-19 Vaccine single-dose (N=156), or a Pfizer-BioNTech COVID-19 vaccine 2-dose series (N=151) at least 12 weeks prior to enrollment and who reported no history of SARS-CoV-2 infection were randomized 1:1:1 to receive a booster dose of one of three vaccines: Moderna COVID-19 Vaccine, Janssen COVID-19 Vaccine, or Pfizer-BioNTech COVID-19 Vaccine. Neutralizing antibody titers, as measured by a pseudovirus neutralization assay using a lentivirus expressing the SARS-CoV-2 Spike protein with D614G mutation, were assessed prior to administration of the booster dose (Day 1) and after the booster dose (Day 15). A booster response to the Pfizer-BioNTech COVID-19 Vaccine was demonstrated regardless of which COVID-19 vaccine was used for primary vaccination. An overall review of adverse reactions reported following the Pfizer-BioNTech COVID-19 vaccine heterologous booster dose did not identify any new safety concerns as compared to adverse reactions reported following the Pfizer-BioNTech COVID-19 vaccine primary vaccination or homologous booster dose (FDA, 2022).

Recent studies indicate that additional doses in people who are moderately or severely immunocompromised are safe and can increase antibody response. Small studies in solid organ transplant recipients in Toulouse, Strasbourg, and Baltimore demonstrate immunogenicity of a 4<sup>th</sup> mRNA dose when administered 1–2 months after the 3<sup>rd</sup> dose (Kamar et al., 2021; Benotmane et al., preprint; Alejo et al., 2021). Multiple studies, including COV-BOOST and the NIH mix-and-match study demonstrated safety and immunogenicity of a booster

dose in the general population when administered at intervals as short as 3 months following a 2-dose primary series (Munro et al., 2021; Atmar et al., preprint). Finally, multiple countries have implemented booster doses at least 3 months after primary vaccination in the general population (e.g., UK, Germany, Netherlands).

### **Risk-Benefit of the COVID-19 vaccine by Pfizer-BioNTech Additional Primary or Booster Vaccination for Individuals Described in the EUI**

The duration of vaccine-induced protection from primary vaccination with COVID-19 vaccines is unknown. Efficacy data from clinical studies of 2-dose primary series supported benefit of the COVID-19 vaccine by Pfizer-BioNTech in preventing severe COVID-19 and supported its FDA approval. Effectiveness of an additional primary dose of the COVID-19 vaccine by Pfizer-BioNTech is inferred from immunogenicity data in immunocompromised adults who received a single additional primary dose. Clinical trials demonstrated that relative vaccine efficacy was 95.3% (95% confidence interval: 89.5%, 98.3%) among persons aged 16 years and older who received a booster dose of the COVID-19 vaccine by Pfizer-BioNTech (administered predominantly between 10–12 months following completion of primary series) in the previous 2 months, compared to those who had only completed two primary doses. Rates of local or systemic adverse events in these trials were similar or lower after a booster dose than after the second primary dose (Perez, 2021; Gruber, 2021). Additionally, the [real world data from the Ministry of Health of Israel](#) that included over 6,300 individuals aged 12–15 years and over 4.1 million individuals 16 years and older who received a booster dose of the COVID-19 vaccine by Pfizer-BioNTech at least 5 months after primary vaccination revealed no new safety concerns.

Effectiveness of a heterologous booster dose of COVID-19 vaccine by Pfizer-BioNTech is inferred from data in adults who received a booster dose following primary vaccination with the Pfizer-BioNTech COVID-19 vaccine or another FDA-authorized COVID-19 vaccine. Available data on the safety or efficacy of a Pfizer-BioNTech vaccine dose after receipt of a non-FDA authorized or approved COVID-19 vaccine are limited. However, based on available information, it appears reasonable to anticipate that known and potential risks of an additional primary dose or a booster dose of the COVID-19 vaccine by Pfizer-BioNTech may be outweighed by its likely benefit to enhance or restore protection by the primary vaccination, which might have waned over time, especially in people with immunocompromising conditions or taking immunosuppressive medications who may require a shorter interval for booster doses.

Refer to the CDC's [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for additional information.

### **Available Alternatives**

Currently, the Pfizer-BioNTech COVID-19 vaccine and Moderna COVID-19 vaccine are the only FDA-approved vaccines for which EUI provide for primary, additional, and/or booster dose administration.

### **Reporting Adverse Event or Medication Errors**

The vaccination provider is responsible for mandatory reporting of the following to the Vaccine Adverse Event Reporting System (VAERS):

- vaccine administration errors whether or not associated with an adverse event,
- serious adverse events (irrespective of attribution to vaccination),
- cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and
- cases of COVID-19 that result in hospitalization or death.

Complete and submit reports to VAERS online at <https://vaers.hhs.gov/reportevent.html>.

For further assistance with reporting to VAERS call 1-800-822-7967.

## References

Alejo J., et al. Antibody Response to a Fourth Dose of a SARS-CoV-2 Vaccine in Solid Organ Transplant Recipients: A Case Series. *Transplantation* 2021;105(21):e280-e281.

Araos R, Jara A. Covid-19 vaccine effectiveness assessment in Chile - 25 October 2021 ([https://cdn.who.int/media/docs/default-source/blue-print/chile\\_rafael-araos\\_who-vr-call\\_25oct2021.pdf?sfvrsn=7a7ca72a\\_7](https://cdn.who.int/media/docs/default-source/blue-print/chile_rafael-araos_who-vr-call_25oct2021.pdf?sfvrsn=7a7ca72a_7))

Atmar, R.L., et al. Heterologous SARS-CoV-2 Booster Vaccinations – Preliminary Report. *medRxiv* Preprint. doi: [10.1101/2021.10.10.21264827](https://doi.org/10.1101/2021.10.10.21264827)

Benotmane, I., Bruel T., Planas, D., Fafi-Kremer, S., Schwarz, O., Caillard, S. A fourth dose of the mRNA-1273 SARS-CoV-2 vaccine improves serum neutralization against the delta variant in kidney transplant recipients. Preprint: <https://www.medrxiv.org/content/10.1101/2021.11.25.21266704v1>

FDA. Fact Sheet for Healthcare Providers Administering Vaccine. Emergency Use Authorization of the Pfizer-BioNTech COVID-19 Vaccine to Prevent Coronavirus Disease 2019, Revised January 3, 2022 <https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/comirnaty-and-pfizer-biontech-covid-19-vaccine>

Gruber, W. BNT162b2 [COMIRNATY® (COVID-19 Vaccine, mRNA)] Booster (Third) Dose. Advisory Committee on Immunization Practices meeting, September 22, 2021: <https://www.cdc.gov/vaccines/acip/meetings/downloads/slides-2021-09-22/02-COVID-Gruber-508.pdf>

Joint Committee on Vaccination and Immunisation (JCVI). JCVI statement regarding a COVID-19 booster vaccine programme for winter 2021 to 2022. London (United Kingdom): Department of Health & Social Care; 14 Sept 2021. <https://www.gov.uk/government/publications/jcvi-statement-september-2021-covid-19-booster-vaccine-programme-for-winter-2021-to-2022/jcvi-statement-regarding-a-covid-19-booster-vaccine-programme-for-winter-2021-to-2022>.

Joint Committee on Vaccination and Immunisation (JCVI) advice on third primary dose vaccination. London (United Kingdom): Department of Health & Social Care; 1 Sept 2021. <https://www.gov.uk/government/publications/third-primary-covid-19-vaccine-dose-for-people-who-are-immunosuppressed-jcvi-advice/joint-committee-on-vaccination-and-immunisation-jcvi-advice-on-third-primary-dose-vaccination>.

Kamar, N., Abravanel, F., Marion O. Assessment of 4 doses of SARS-CoV-2 messenger RNA-based vaccine in recipients of a solid organ transplant. *Infection Diseases* 2021;4(11):e2136030.

Meeting of the Strategic Advisory Group of Experts on Immunization (SAGE) – 5 October 2021. Geneva: World Health Organization; 2021. Kaslow DC. Policy question: Is a 3<sup>rd</sup> dose needed for older persons who have received 2 doses of inactivated COVID-19 vaccines? [https://www.who.int/news-room/events/detail/2021/10/04/default-calendar/sage\\_meeting\\_october\\_2021](https://www.who.int/news-room/events/detail/2021/10/04/default-calendar/sage_meeting_october_2021).

Moghnieh R, Mekdashi R, El-Hassan S, et al. Immunogenicity and reactogenicity of BNT162b2 booster in BBIBP-CorV-vaccinated individuals compared with homologous BNT162b2 vaccination: Results of a pilot prospective cohort study from Lebanon. *Vaccine* 2021; 139:6713-9.

Munro, Alasdair PS, et al. "Safety and immunogenicity of seven COVID-19 vaccines as a third dose (booster) following two doses of ChAdOx1 nCov-19 or BNT162b2 in the UK (COV-BOOST): a blinded, multicentre, randomised, controlled, phase 2 trial." *The Lancet* (2021).

National Advisory Committee on Vaccination. An Advisory Committee Statement: Guidance on booster COVID-19 vaccine doses in Canada. Public Health Agency of Canada; 3 Dec 2021.

<https://www.canada.ca/content/dam/phac-aspc/documents/services/immunization/national-advisory-committee-on-immunization-naci/guidance-booster-covid-19-vaccine-doses/guidance-booster-covid-19-vaccine-doses.pdf>

Patamatamkul S, Thammawat S, Buranrat B. Induction of robust neutralizing antibodies against the COVID-19 Delta variant with ChAdOx1 nCoV-19 or BNT162b2 as a booster following a primary vaccination series with CoronaVac. medRxiv 2021.09.25.21264099; doi: <https://doi.org/10.1101/2021.09.25.21264099>.

Perez, J. Efficacy and safety of BNT162b2 booster – C4591031 2 month interim analysis. Advisory Committee on Immunization Practices meeting, November 19, 2021:

<https://www.cdc.gov/vaccines/acip/meetings/downloads/slides-2021-11-19/02-COVID-Perez-508.pdf>

WHO Strategic Advisory Group of Experts on Immunization (SAGE). Interim recommendations for an extended primary series with an additional vaccine dose for COVID-19 vaccination in immunocompromised persons; updated 26 October 2021. Geneva: World Health Organization; 2021.

[https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE\\_recommendation-immunocompromised-persons](https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE_recommendation-immunocompromised-persons).

WHO Strategic Advisory Group of Experts on Immunization (SAGE). Interim recommendations for use of the inactivated COVID-19 vaccine, CoronaVac, developed by Sinovac Interim guidance; updated 21 October 2021.

Geneva: World Health Organization; 2021. [https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE\\_recommendation-Sinovac-CoronaVac-2021.1](https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE_recommendation-Sinovac-CoronaVac-2021.1).

WHO Strategic Advisory Group of Experts on Immunization (SAGE). Interim recommendations for use of the inactivated COVID-19 vaccine BIBP developed by China National Biotec Group (CNBG), Sinopharm; updated 28 October 2021. Geneva: World Health Organization; 2021. <https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE-recommendation-BIBP>.