



State of Missouri regional COVID-19 hospitalized cases model

October 14, 2020

Multiple data points inform Missouri's COVID-19 response


- Syndromic surveillance
- Healthcare system capacity (bed, PPE, and staff availability)
- Testing
- COVID-19 cases and deaths
- Economic and social impact
- Insights from U.S. states, nationally, and other countries
- Evidence from scientific literature
- Mathematical disease modelling

Our model estimates possible outcomes based on currently available information

What does the model tell us	What does it not tell us
Range of plausible outcomes based on our current knowledge of COVID-19 in Missouri	What will happen in the future
Approximate date and magnitude of peak/s based on current understanding of policy interventions and human behavior and assumptions about future interventions	Date and magnitude of peak/s if there are major changes in planned policy interventions and human behavior
Approximate estimate of effective transmission rate across a region	Exact transmission rate in all parts of a region – there may be areas of higher and lower transmission within the region
Projected hospitalizations for regions in MO with sufficient data, i.e. Kansas City Area, Central, St. Louis Area, Southeast and Southwest	Projected hospitalizations in regions where daily COVID-19 hospitalizations are fewer than 15 because insufficient cases

The ability to forecast depends on the quality and availability of data. For a new disease such as COVID-19, much remains uncertain.

- “ R_e ” rates near or above 1 in nearly every region means the disease is spreading statewide



* Data date range: 10/06/20 - 10/13/202

Central (Region F)




Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	2.3		% ICU Beds Occupied	67%	
Last Week	1.17		% ICU Beds Occupied C19	16%	
Current Week	1.16		% ICU Beds Free	33%	
WoW % Change	-0.6%				
			% Ventilators in use	37%	
			% Ventilators available	63%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Central Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Greater Kansas City Area (Region A)




Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	2.8		% ICU Beds Occupied	70%	
Last Week	1.11		% ICU Beds Occupied C19	12%	
Current Week	1.13		% ICU Beds Free	30%	
WoW % Change	1.4%				
			% Ventilators in use	25%	
			% Ventilators available	75%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Kansas City Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Northeast (Region B)




Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	N/A		% ICU Beds Occupied	68%	
Last Week	0.81		% ICU Beds Occupied C19	77%	
Current Week	0.87		% ICU Beds Free	32%	
WoW % Change	6.8%				
			% Ventilators in use	9%	
			% Ventilators available	91%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Northeast Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Northwest (Region H)




Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	1.24		% ICU Beds Occupied	88%	
Last Week	1.255		% ICU Beds Occupied C19	34%	
Current Week	1.255		% ICU Beds Free	13%	
WoW % Change	0.0%				
			% Ventilators in use	19%	
			% Ventilators available	81%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Northwest Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Southeast / Cape Girardeau (Region E)



Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	2.61		% ICU Beds Occupied	63%	
Last Week	1.11		% ICU Beds Occupied C19	22%	
Current Week	1.15		% ICU Beds Free	37%	
WoW % Change	3.7%				
			% Ventilators in use	34%	
			% Ventilators available	66%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Southeast Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Southwest / Springfield (Regions D,G, I)



Reproductive Rate		
Pre-intervention	2.36	
Last Week	1.11	
Current Week	1.10	
WoW % Change	-0.3%	


Bed / Ventilator Availability		
% ICU Beds Occupied	70%	
% ICU Beds Occupied C19	23%	
% ICU Beds Free	30%	
% Ventilators in use	27%	
% Ventilators available	73%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients

Base Case Southwest Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Greater St Louis Area (Region C)




Reproductive Rate			Bed / Ventilator Availability		
Pre-intervention	3.39		% ICU Beds Occupied	71%	
Last Week	1.03		% ICU Beds Occupied C19	14%	
Current Week	1.03		% ICU Beds Free	29%	
WoW % Change	0.1%				
			% Ventilators in use	40%	
			% Ventilators available	60%	

* % of ICU beds occupied by COVID-19 PUI/Confirmed patients


Base Case St. Louis Region

Model Scenario: Base Case, From Date: Mar 1, 2020, To Date: 10/28/20 1:00 AM, + 2 more

Hospitalizations



Version 1.0, As of June 3, 2020




Missouri's Regional COVID-19 Hospitalized Cases Model: Overview and Frequently Asked Questions

Model Overview

One of the many data analyses that inform Missouri's COVID-19 response is a regional model of hospitalized COVID-19 cases that the State of Missouri developed in partnership with the Washington University in St. Louis and Missouri Hospital Association.

Missouri's model uses a standard SEIR (susceptible, exposed, infectious, recovered) compartmental structure that is based upon a tool called LEMMA (Local Epidemiological Modeling for Management & Action), which was developed by experts from UMass Amherst, UC Berkeley, UCSF, and WUStL. The model focuses on COVID-19 hospitalized cases to directly address the question of hospital capacity and provide a more accurate picture of COVID-19's impact on the healthcare system.



```

graph LR
    Susceptible --> Exposed
    Exposed --> Infectious
    Infectious --> Removed
    Infectious --> Hospitalized
    Hospitalized --> Removed
    Hospitalized --> Floor
    Hospitalized --> ICU
    Hospitalized --> Vent
  
```

To help inform decisions at the regional and local level, each region is modeled separately using the latest local data, including COVID-19 confirmed and suspected hospitalizations, population, policy interventions, and average hospital length of stay.

General FAQs

Why are regional models of COVID-19 important?

When new diseases such as COVID-19 emerge, there is much uncertainty about how best to control the epidemic. Decision makers must make the best possible decisions with the available information at hand.

Mathematical models are commonly used to make projections of how infectious diseases might impact key outcomes such as hospitalized cases or deaths. Today, there are many sophisticated models of COVID-19 that make global or national projections (e.g., Imperial College, Harvard, [UW-Madison](#)). However, these generally do not incorporate key local or regional inputs, such as variations in local population demographics, healthcare system

capacity, and the effectiveness of interventions. The model developed by the Missouri Hospital Association and Washington University in St. Louis addresses these limitations by providing a more accurate picture of COVID-19's impact on the healthcare system.

Low levels of daily COVID-19 hospitalizations in the Northeast and Northwest regions limit the ability to generate projections for these regions. In particular, the numbers of hospitalized cases have been so low that

projections for these regions are not reliable. The model focuses on COVID-19 hospitalized cases to directly address the question of hospital capacity and provide a more accurate picture of COVID-19's impact on the healthcare system.

Missouri's model uses a standard SEIR (susceptible, exposed, infectious, recovered) compartmental structure that is based upon a tool called LEMMA (Local Epidemiological Modeling for Management & Action), which was developed by experts from UMass Amherst, UC Berkeley, UCSF, and WUStL. The model focuses on COVID-19 hospitalized cases to directly address the question of hospital capacity and provide a more accurate picture of COVID-19's impact on the healthcare system.

Regional COVID-19 transmission models help inform local policy, public health, and business decisions

- Mathematical models are commonly used to make projections of infectious disease epidemics (e.g., tuberculosis, HIV)
- Many sophisticated models on COVID-19 make global or national projections (e.g., Imperial College, Harvard, IHME)
- However, these generally do not incorporate critical local or regional inputs, such as:
 - Variations in local population size and age structure
 - Date and nature of social distancing and other policies
- Regional projections are important because:
 - Regional epidemics may differ markedly from the national average
 - Policy response occurs at state, county, and municipal levels

State of MO, WUSTL, and MHA have developed a regional model of hospitalized COVID-19 cases

- **Standard SEIR model that combines universal characteristics of COVID-19 infection (e.g., transmission parameters) with local inputs to support regional decision making**
 - Mathematical model developed by experts from UMass Amherst, UC Berkeley, UCSF, and WUSTL
 - Uses a statistical approach that adjusts underlying parameters as new data are observed
- **Customized using the latest local data from Missouri's emergency response regions, including:**
 - COVID-19 positives and PUIs
 - Population and age structure
 - Policy interventions
 - Avg. hospital length of stay
- **Projects COVID-19 hospitalized cases** to directly address the question of hospital capacity and provide a more accurate picture on COVID-19's impact on the healthcare system

Model Structure (SEIR)

