

Draft LOC Algorithm 2.2

Proposed Level of Care Threshold: 18 points

When testing the algorithm, all indicators should be looked at individually to ensure the highest score is being given to the participant in the category. For example, in the behavioral section an individual may only score 1 in wandering but may score a 3 in verbal abuse. Using the higher score in verbal abuse will give the participant 6 points instead of the 3 they would have scored for wandering.

Behavioral: Repeated behavioral challenges that affect a participant's ability to function in the community.

- E3a - Wandering
- E3c - Physical Abuse
- E3d - Socially Inappropriate / Disruptive
- E3e - Inappropriate Public Sexual Behavior
- E3f - Resists Care
- J3g - Abnormal Thought Process
- J3h - Delusions
- J3i - Hallucinations
- N7b - Mental Condition

If (N7b=1)
OR (E3a =1)
OR (E3c=1)
OR (E3d=1)
OR (E3e=1)
OR (E3f=1)
OR (J3g=1)
OR (J3h=1)
OR (J3i=1)
Then LOC = 3

If (N7b=2 or N7b=3)
OR (E3a=2 or E3a=3)
OR (E3c=2 or E3c=3)
OR (E3d=2 or E3d=3)
OR (E3e=2 or E3e=3)
OR (E3f=2 or E3f=3)
OR (J3g=2 or J3g=3 or J3g=4)
OR (J3h=2 or J3h=3 or J3h=4)
OR (J3i=2 or J3i=3 or J3i=4)
Then LOC = 6

If (N7b=2 or N7b=3)
AND
(E3a=3)
OR (E3c=3)

Draft LOC Algorithm 2.2

OR (E3d=3)

OR (E3e=3)

OR (E3f=3)

OR (J3g=3 **or** J3g=4)

OR (J3h=3 **or** J3h=4)

OR (J3i=3 **or** J3i=4)

Then LOC = 9

Cognition: Assesses a participant's performance in remembering, making decisions, organizing daily self-care activities, as well as understanding others and making self-understood.

If a participant scores C1=5 (no discernable consciousness, coma, as outlined in the InterRAI) the participant is presumed to require nursing facility LOC; this is defined as a TRIGGER.

- C1 - Cognitive Skills
- C2a - Short Term Memory
- C2b - Procedural Memory
- C2c - Situational Memory
- C3c - Mental Function
- D1 - Making Self Understood
- D2 - Ability to Understand Others

If (C1=1 **or** C1=2)

AND

(C2a=1)

OR (C2b=1)

OR (C2c=1)

OR (C3c=1 **or** C3c=2)

OR (D1=2 **or** D1=3 **or** D1=4)

OR (D2=2 **or** D2=3 **or** D2=4)

Then LOC = 3

If (C1=3)

AND

(C2a=1)

OR (C2b=1)

OR (C2c=1)

OR (C3c=1 **or** C3c=2)

OR (D1=3)

OR (D2=3)

Then LOC = 6

If (C1=3)

AND

(D1=4)

OR (D2=4)

Draft LOC Algorithm 2.2

OR

(C1=4)

Then LOC = 9

If (C1=5)

Then LOC = 18 *TRIGGER

Activities of Daily Living (ADLs)

Activities of Daily Living or ADLs refer to the fundamental activities an individual does on a day-to-day basis necessary for independent living. The scoring for Mobility, Eating and Toileting reflects 0, 3, 6 or 9. The scoring for Bathing and Dressing reflects 0, 3 or 6. The scoring for Missouri is based upon the foundation of the Hierarchy of ADLs, (InterRAI).

Mobility: Amount of assistance needed to move from one place or position to another.

G2f - Locomotion
G2i - Bed Mobility
G3a - Primary mode of locomotion

If (G2f=3 or G2f=4)

OR

(G2i=3 or G2i=4)

Then LOC = 3

If (G2f=5)

OR

(G2i=5 or G2i=6)

Then LOC = 6

If (G3a=3)

OR

(G2f=6)

Then LOC = 18 *TRIGGER

*If a participant scores G3a=3 (primary mode of locomotion, bedbound, as outlined in the InterRAI) the participant is presumed to require nursing facility LOC; this is defined as a TRIGGER.

Eating: Amount of assistance needed to eat and drink, special nutritional requirements and mode of nutrition.

G2j - Eating
K2e - Therapeutic Diet

Draft LOC Algorithm 2.2

If (G2j=1 or G2j=2 or G2j=3)
OR
(K2e=1)
Then LOC = 3

If (G2j=4)
Then LOC = 6

If (G2j=5)
Then LOC = 9

If (G2j=6)
Then LOC = 18 *TRIGGER

*If a participant scores G2j=6 (eating, total dependence, as outlined in the InterRAI) the participant is presumed to require nursing facility LOC; this is defined as a TRIGGER.

Toileting: Amount of assistance needed to use the toilet room (or commode, bedpan, urinal), transfers on/off toilet, cleanses, changes (incontinence), pad, manages ostomy or catheter, and adjusts clothes.

G2g - Transfer Toilet
G2h - Toilet Use

If (G2g=3 or G2g=4)
OR
(G2h=3 or G2h=4)
Then LOC = 3

If (G2g=5)
OR
(G2h=5)
Then LOC = 6

If (G2g=6)
OR
(G2h=6)
Then LOC = 9

Bathing: Amount of assistance needed for a full-body bath/shower.

G2a - Bathing

If (G2a=3 or G2a=4)

Draft LOC Algorithm 2.2

Then LOC = 3

If (G2a=5 or G2a=6)
Then LOC = 6

Dressing and Grooming: Amount of assistance needed to dress and undress (street clothes, underwear) including prostheses, orthotics, fasteners, pullovers, etc.

G2b - Personal Hygiene
G2c - Dressing Upper Body
G2d - Dressing Lower Body

If (G2b=3 or G2b=4)
OR
(G2c =3 or G2c=4)
OR
(G2d=3 or G2d=4)
Then LOC = 3

If (G2b=5 or G2b=6)
OR
(G2c=5 or G2c=6)
OR
(G2d=5 or G2d=6)
Then LOC = 6

Rehabilitation: The restoration of a former or normal state of health through medically-ordered therapeutic services either directly provided by or under the supervision of a qualified professional.

N3ea - PT
N3fa - OT
N3ga - Speech
N3ia - Cardiac Rehab

If (N3ea = 1)
OR
(N3fa = 1)
OR
(N3ga = 1)
OR
(N3ia = 1)
Then LOC = 3

If (N3ea = 2 or N3ea = 3)

Draft LOC Algorithm 2.2

OR

(N3fa = 2 **or** N3fa = 3)

OR

(N3ga = 2 **or** N3ga = 3)

OR

(N3ia = 2 **or** N3ia = 3)

Then LOC = 6

If (N3ea = 4 **or** N3ea = 5 **or** N3ea = 6 **or** N3ea = 7)

OR

(N3fa = 4 **or** N3fa = 5 **or** N3fa = 6 **or** N3fa = 7)

OR

(N3ga = 4 **or** N3ga = 5 **or** N3ga = 6 **or** N3ga = 7)

OR

(N3ia = 4 **or** N3ia = 5 **or** N3ia = 6 **or** N3ia = 7)

Then LOC = 9

Treatments: Physician ordered medical care or management that requires additional hands on assistance. The scoring for treatments will be 0 or 6. Participants with the identified treatments will receive 6 points.

H1	-	Bladder Continence
H2	-	Urinary Collection Device
H3	-	Bowel Continence
K3	-	Mode of Nutrition
L1	-	Pressure Ulcer Severity
L3	-	Presence of Skin Ulcer
L4	-	Major Skin Problems
L5	-	Skin Tears or Cuts
N2g	-	Suctioning
N2h	-	Tracheostomy Care
N2j	-	Ventilator or Respirator
N2k	-	Wound Care

If (H1=1)

OR

(H2=1 **or** H2=2 **or** H2=3)

OR

(H3=1)

OR

(K3=5 **or** K3=6 **or** K3=7 **or** K3=8)

OR

(N2g=1 **or** N2g=2 **or** N2g=3 **or** N2g=4)

OR

(N2h=1 **or** N2h=2 **or** N2h=3 **or** N2h=4)

OR

Draft LOC Algorithm 2.2

(N2j=1 or N2j=2 or N2j=3 or N2j=4)

OR

(N2k=1 or N2k=2 or N2k=3 or N2k=4)

AND

(L1=2 or L1=3 or L1=4 or L1=5 or L1=6)

OR

(L3=1)

OR

(L4=1)

OR

(L5=1)

Then LOC = 6

Instrumental Activities of Daily Living (IADL): Instrumental activities of daily living are the skills and abilities needed to perform certain day-to-day tasks associated with an independent lifestyle.

Managing Medications: Amount of assistance needed to safely manage medications based on the participant's capacity to complete the task.

- B4c - Mental Health Residence
- B4d - Psychiatric Hospital or Unit
- B4e - Settings for Persons with Intellectual Disability
- C1 - Cognitive Skills
- C2b - Procedural Memory
- C3c - Mental Function
- G1d - Managing Medications

If (G1d=2)

AND

(B4c=1)

OR

(B4d=1)

OR

(B4e=1)

OR

(C1=2 or C1=3 or C1=4 or C1=5)

OR

(C2b=1)

OR

(C3c=1 or C3c=2)

OR

(G1d=3 or G1d=4)

Then LOC = 3

If (G1d=5 or G1d=6)

Draft LOC Algorithm 2.2

Then LOC = 6

Meal Prep: Amount of assistance needed to prepare a meal based on the participant's capacity to complete the task.

G1a - Meal Prep

If (G1a=3 or G1a=4)
Then LOC= 3

If (G1a=5 or G1a=6)
Then LOC = 6

Safety: Participant is at risk of injury due to visual impairment, unsteady gait, and/or falls.

* Age 75 or greater – add 3 additional points to the participant's highest score received in the safety category.

- B4a - Long Term Care Facility
- B4b - RCF/ALF
- B4c - Mental Health Residence
- B4d - Psychiatric Hospital or Unit
- B4e - Settings for Persons with Intellectual Disability
- D4 - Vision
- J1 - Falls
- J3a - Problem frequency to move to standing position
- J3b - Problem frequency to turn to face opposite direction
- J3c - Problem frequency for dizziness
- J3d - Problem frequency for unsteady gait

If (B4a=1)
OR
(B4b=1)
OR
(B4c=1)
OR
(B4d=1)
OR
(B4e=1)
OR
(D4=3)
OR
(J1=1 or J1=2 or J1=3)
OR
(J3a=2 or J3a=3 or J3a=4)

Draft LOC Algorithm 2.2

OR
(J3b=2 **or** J3b=3 **or** J3b=4)

OR
(J3c=2 **or** J3c=3 **or** J3c=4)

OR
(J3d=2 **or** J3d=3 **or** J3d=4)
Then LOC = 3 *

If (D4=4)
OR (J1=1 **or** J1=2 **or** J1=3)
AND
(J3a=2 **or** J3a=3 **or** J3a=4)
OR
(J3b=2 **or** J3b=3 **or** J3b=4)
OR
(J3c=2 **or** J3c=3 **or** J3c=4)
OR
(J3d=2 **or** J3d=3 **or** J3d=4)
Then LOC = 6 *

If Max score of 6 prior the addition of age points
AND
75 years of age or older
Then LOC = 18 TRIGGER *

*If a participant received the maximum score in safety and is 75 years of age or older the participant is presumed to require nursing facility LOC; this is defined as a TRIGGER.